

Introduction to General Relativity

Janos Polonyi

University of Strasbourg, Strasbourg, France

(Dated: October 13, 2017)

Contents

I. Introduction	4
A. Equivalence principle	4
B. Uniformly accelerating observer	6
C. Static gravitational field	7
D. Classical field theories	8
II. Gauge theories	9
A. Global symmetries	10
B. Local symmetries	10
C. Covariant derivative	13
D. Parallel transport	15
E. Field strength tensor	16
III. Gravity	18
A. Classical field theory on curved space-time	18
B. Geometry	21
C. Gauge group	22
1. Space-time diffeomorphism	23
2. Internal Poincaré group	24
D. Gauge theory of diffeomorphism	25
1. Covariant derivative	25
2. Lie derivative	28
3. Field strength tensor	29
E. Metric admissibility	30
F. Dynamics	35

IV. Coupling to matter	37
A. Point particle in an external gravitational field	37
1. Equivalence Principle	37
2. Spin precession	38
3. Variational equation of motion	39
4. Geodesic deviation	40
5. Newtonian limit	41
B. Interacting matter-gravity system	42
1. Point particle	42
2. Ideal fluid	43
3. Classical fields	44
V. Gravitational radiation	44
A. Linearization	45
B. Wave equation	46
C. Plane-waves	47
D. Polarization	47
VI. Schwarzschild solution	49
A. Metric	49
B. Geodesics	53
C. Space-like hyper-surfaces	55
D. Around the Schwarzschild-horizon	56
1. Falling through the horizon	57
2. Stretching the horizon	57
3. Crossing the horizon	59
4. Reduplication of space-time	59
5. Causal structure	62
VII. Homogeneous and isotropic cosmology	63
A. Maximally symmetric spaces	64
B. Robertson-Walker metric	65
C. Equation of motion	68
D. Frequency shifts	71

E. Particle horizon	73
F. Evolution of the universe	75
A. Classical Field theory	77
1. Variational principle	77
a. Single point on the real axis	78
b. Non-relativistic point particle	78
c. Scalar field	80
2. Noether theorem	81
a. Point particle	82
b. Internal symmetries	83
c. External Symmetries	84
B. Parallel transport along a path	86
C. Higgs mechanism in scalar electrodynamics	87
1. Spontaneous symmetry breaking	87
2. Goldstone theorem	88
3. Higgs mechanism	89
D. Scalar particle on a homogeneous non-Abelian magnetic field	90
E. Gauge theory of the Poincaré group	91
1. Covariant derivatives	92
a. Vierbein	92
b. Holonomic and anholonomic vector fields	93
c. Local Lorentz transformations	94
d. Local translations	95
e. Full local Poincare group	96
f. World vectors and tensors	96
2. Field strength tensors	97
3. Variational equations	98
a. Variation of ω	99
b. Variation of e	101
c. Affine connection for world vectors	103

I. INTRODUCTION

A distinguishing feature of gravitation compared with the other known interactions in Physics is that it touches upon the geometry of space and time. Correspondingly, the traditional approach to General Relativity is based on differential geometry. But there is another issue, locality which historically came up in connection with gravity first with fully recognized importance. Locality is evoked at two different stages in General Relativity. One is the independence of the equations of the theory from the choice of space-time coordinates. This renders the theory invariant under local reparametrization of the coordinate system. The equivalence principle, the existence of coordinate systems where gravitational forces disappear at a given point is another indication that locality plays more important role than thought before.

But locality actually represents a bridge to the other interactions. In fact, the interplay of locality and Special Relativity leads to gauge theories, a formal structure, common in all interactions known to us, gravity included. Inspired by this common structure General relativity is introduced below as a classical gauge field theory. After that a brief description of two applications is given, namely the Schwarzschild geometry and the Robertson-Walker geometry based cosmology.

Two appendices contain some complementary material, a short introduction into the formalism of classical field theory and the presentation of General Relativity as the gauge theory of the Poincaré group.

A. Equivalence principle

The unique status of gravitational forces among possible interactions is that surprising equivalence of the inertial mass appearing in Newton's third law in mechanics, m_{in} , and the coupling strength to a gravitational potential $\check{\mathbf{J}}_{gr}$, the gravitational mass m_{gr} ,

$$m_{in}\mathbf{a} = -m_{gr}\nabla\check{\mathbf{J}}_{gr}(\mathbf{x}). \quad (1)$$

As a result, the trajectory of a point particle is independent of its mass within Newton's theory. Lorand Eötvös' measurement of the late nineteenth century and the improved versions performed later show the equivalence of gravitational and inertial masses with convincingly high accuracy. The Weak Equivalence Principle states that the world line of a small, free falling body is independent of its composition or structure.

The equivalence,

$$m_{in} = m_{gr} = m, \quad (2)$$

can be interpreted as a sign that the gravitational force has the same origin as the inertial forces, arising from using an accelerating coordinate system. In fact, assuming the presence of an external force, \mathbf{F}_{ext} , acting on the particle beyond the gravitational one the equation of motion reads as

$$m[\mathbf{a} + \nabla \check{\mathbf{J}}_{gr}(\mathbf{x})] = \mathbf{F}. \quad (3)$$

For instance, a homogeneous gravitational field, $\check{\mathbf{J}}_{gr}(\mathbf{x}) = gz$, can be eliminated by means of an accelerating coordinate system, $z \rightarrow z - gt^2/2$, one is in a levitation, weightless state in the falling elevator. Such an equivalence of inertial and gravitational forces is a local phenomenon, an inhomogeneous, time dependent gravitational potential can be eliminated at a given space-time point only by means of a suitably chosen space-time coordinates. In other words, the usual dynamics and symmetries, predicted by Special Relativity can be recovered in the absence of non-gravitational forces at any given point in space-time by means of a well chosen coordinate system. The generalization of this statement for any interaction is the Strong Equivalence Principle, stating that for an observer in free fall in a gravitational field the results of all local experiments are independent of the strength of the gravitational field. Since all fundamental laws in physics, this, too, holds locally only.

One can rephrase the Strong Equivalence Principle by saying that in a small enough region of space-time the laws of physics reduce to those of Special Relativity and it is impossible to detect the presence of a gravitational field. In other words, the space-time and the physical laws can be made locally Lorentz-invariant. The mathematical details of locally observed Lorentz invariance will be spelled out below as a gauge symmetry. Yet another form of the Equivalence Principle is the equivalence of inertial forces which arise from the use of some coordinate system in space-time and the gravitational interaction.

Note that the gravitational forces are assumed to be originating from a simple scalar potential in this discussion, a restriction to be released later. But the principle is not flawless and has a limited validity, for instance quantum effects, related to the spin of the particles seem to represent an $\mathcal{O}(\hbar)$ violation. But quantum effects will be ignored in the rest of this notes. There are several, slightly different versions of the Equivalence Principle in the literature which confuse the picture even more. One may say that the Equivalence Principle is a rough guidance for our intuition which comes from the pre-gauge theory era of physics. Its simpler form and its limitation can easily be found when gravity is considered as a gauge theory.

The equivalence of the inertial and the gravitating mass, (2), makes that the only characteristic classical quantity of the point particle, its mass, drops out from the gravitational dynamics. But

what determines then the particle trajectory if not its mass? It has been suspected ever since the construction of Riemann-geometry that the gravitational force ought to be related to the curvature because it determines the geometry. In fact, consider the free fall in the (x, y) plane, $x(t) = vt$, $y(t) = y_0 - gt^2/2$. The curvature of the trajectory, the radius of the sphere touching the trajectory, $R = 1/(d^2y/dx^2) = -v^2/g$ depends on the gravitational constant, g . In other words, the gravitational force makes the particle to follow a trajectory with a given curvature. But the curvature contains the initial velocity, v , as well, hence it is not of purely geometric origin. It was Einstein's radically new point of view that the trajectory should be viewed in the space-time rather than in the space only where the curvature becomes independent of the initial conditions and can be assigned to geometry alone.

B. Uniformly accelerating observer

If inertial and gravitational forces are identical then a uniformly accelerating observer should experience a homogeneous gravitational potential and it is instructive to look more closely in this case. Consider the first motion in the spatial x direction where the world line $x^\mu(s)$,

$$x^\mu = (t, x) = \frac{1}{a}(\sinh a\tau, \rho - 1 + \cosh a\tau), \quad (4)$$

s being the invariant length, gives rise the four-velocity $u^\mu(s) = \frac{d}{ds}x^\mu(s) = \dot{x}^\mu(s)$ which satisfies $u^2 = 1$ and the four acceleration $a^\mu(s) = \ddot{x}^\mu(s)$ with constant invariant length, $a^2 = -a_0^2$. The four-velocity, written as $u^\mu = (\cosh f(s), \sinh f(s), 0, 0)$, gives the desired acceleration with the choice $f(s) = as$. An integration of the velocity produces the world line

$$x^\mu = \frac{1}{a}(\sinh as, \cosh as, 0, 0), \quad (5)$$

a hyperbole with positive value of the x coordinate. It is advantageous to introduce a coordinate system $(t, x) \rightarrow (\eta, \rho)$, where the hyperboles are labeled by the spatial coordinate ρ and η is proportional to the proper time,

$$x^\mu = \rho(\sinh \eta, \cosh \eta, 0, 0), \quad (6)$$

where the proper time and the acceleration of the world lines are $s = \eta/\rho$ and $a = 1/\rho$, respectively and the invariant distance can be written as

$$ds^2 = \rho^2 d\eta^2 - d\rho^2. \quad (7)$$

The following features of the Rindler geometry are noteworthy:

1. The Rindler space covers the part $x \geq 0$ of the whole Minkowski geometry,
2. The geometry, described by the metric (7) is flat, it is a reparametrization of the Minkowski space-time.
3. The acceleration along a given world line is constant but different world lines display different acceleration hence the gravitational field is static but spatially inhomogeneous.
4. There are gravitational effect even in flat space-time.
5. There is no contradiction with the Equivalence Principle because the accelerating world lines fill up a part of the space-time only and no statement is made about the flatness of the rest.
6. Gravitational forces seem to generate space-time dependent metric tensor.
7. Gravitational forces seem to produce singularity whose more precise nature, namely whether it is a coordinate singularity due to the wrong choice of coordinates or it is a real singularity, reflected by physical quantities, remains to be clarified.
8. Gravitational forces seem to make a part of the space-time inaccessible, they can generate a horizon.

C. Static gravitational field

We have seen a distinguishing feature of gravitational forces, the equivalence principle above. We turn now to the similarity between static gravitational and electric forces, by comparing the Coulomb force

$$\mathbf{F}_C = \mathbf{r} \frac{e_1 e_2}{r^3}, \quad (8)$$

and Newton's gravitational law,

$$\mathbf{F}_g = -\mathbf{r} \frac{\tilde{m}_1 \tilde{m}_2}{r^3}, \quad (9)$$

where e denote the electric charge and $\tilde{m} = m\sqrt{G}$, G being Newton's gravitational constant.

Despite their formal similarity the static electric and weak gravitational forces differ on two counts. One difference is that the gravitational charge, the mass can not be negative as opposed to the electric charge. This eliminates the possibility of screening of gravitational interaction. As a result, the gravitational forces remain long range like the unscreened Coulomb force and lead to

instabilities and a number of interesting differences between the rules of statistical physics with and without gravitational forces. The other difference appears when the charges are not static but accelerate. The resulting electro-dynamical and gravitational radiation are very different from each other.

The dynamical degree of freedom representing the electromagnetic interaction is the vector potential $A_\mu(x)$. The space-time geometry of classical General Relativity is characterized by the invariant length, defined by the metric tensor, $g_{\mu\nu}(x)$ introduced below. This tensor field can be considered as the dynamical degrees of freedom, responsible of gravitational interaction in classical physics.

The vector potential and the metric tensor describe elementary particles with spin one, $S = 1$, and spin two, $S = 2$, respectively. It is now understood that both the absence of gravitational attraction and the complicated structure of gravitational radiation, compared to classical electro-dynamics arise from the different spin of the carriers of the interaction.

An unifying concept, local gauge symmetries, streamlines the construction of classical theories for fields with non-vanishing spin. The minimalistic version of a gauge theory, the Yang-Mills model will be introduced in the next Chapter.

D. Classical field theories

The need of classical fields appear as a way around an unexpected problem in Special Relativity. The non-relativistic equation of motion for a system of N point particles, interacting by instantaneous, action-at-a-distance force is

$$m_a \frac{d^2 \mathbf{x}_a(t)}{dt^2} = \mathbf{F}_a(\mathbf{x}_1(t), \dots, \mathbf{x}_N(t)), \quad (10)$$

where $a = 1, \dots, N$, with initial conditions imposed at $t = t_i$. The relativistically covariant extension of such equations is given for the world lines, $x^\mu(s)$, parametrized by the invariant length s , called proper time, and read

$$m_a \ddot{x}_a^\mu(s_a) = F_a^\mu(x_1(s_1), \dots, x_N(s_N)), \quad (11)$$

where the dot stands for the derivation with respect to the proper time, the proper time of the particles is chosen in such a manner on the right hand side of the equation of motion that $x_a^0(s_a) = x_b^0(s_b)$ and the initial conditions are imposed at $x_a^0(s_i) = t_i$. The problem comes from the fact that the four velocity preserves its length, $\dot{x}_a^2(s) = 1$ and the derivative with respect to the proper

time gives the constraint $0 = \dot{x}_a \ddot{x}_a = \dot{x}_a F_a$ for each world line. One can show that there is no covariant function F_a^μ which satisfies this constraint.

The origin of the problem is that the instantaneous action-at-a-distance interaction requires propagation of signals with infinite velocities which is excluded by Special Relativity. The solution is to represent the interaction by means of a field variable, denoted by $\phi(\mathbf{x})$ for the sake of a simple example, a dynamical degrees of freedom at each space point. To make up the propagation of a signal we couple $\phi(\mathbf{x})$ to the fields of the neighboring points in a relativistically invariant manner. The local dynamics of a field is usually defined by means of its Lagrangian, cf. Appendix A. The simplest form, motivated by the Landau-Ginsburg double expansion is

$$L = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{m^2 c^2}{2\hbar^2} \phi^2, \quad (12)$$

where $\partial_\mu = \partial/\partial x^\mu$, and $\lambda_C = \frac{\hbar}{mc}$ is the Compton wave-length of a particle of mass m , describes a harmonic system. The normal modes, plane waves $\phi(x) = \phi_0 e^{-\frac{i}{\hbar} p x}$, $p = (E/c, \mathbf{p})$ have the dispersion relation, energy-momentum function $E = \pm E_{\mathbf{p}}$ with

$$E_{\mathbf{p}} = c \sqrt{m^2 c^2 + \mathbf{p}^2}. \quad (13)$$

Finally one couples the particles to the field and the result is a system of particles, where the retarded interaction obeys the relativistic symmetries and propagates with velocity $v \leq c$. The field, representing the interaction is usually obeying anharmonic dynamics and its Lagrangian contains higher than quadratic terms in the field.

II. GAUGE THEORIES

We know four fundamental interactions in Nature, the gravitational, the electric, the weak and the strong forces. The last three appears in quantum version but we have so far no experimental evidence whatsoever indicating propagating classical gravitational field or any quantum effect in gravity. The remarkable fact is that all four interactions are described by the same type of models, by gauge field theories. These theories express a central assumption of physics in a distinguished manner, namely that the fundamental laws are local in the space-time. Non-gravitational gauge theories are introduced in this section in flat space-time.

A. Global symmetries

A transformation of the field configuration, $\phi^a(x) \rightarrow \phi'^a(x')$, where $x \rightarrow x'(x)$ is a symmetry if it changes the action $S[\phi]$ at most by total derivative,

$$S[\phi] \rightarrow S[\phi'] = S[\phi] + \int d^4x \partial_\mu \Lambda^\mu(x) \quad (14)$$

because such transformations leave the variational equations of motion unchanged.

It is advantageous to distinguish two different spaces in field theory. A field configuration of an n -component real field, $\phi(x) : E \rightarrow I$, is a mapping of the external space into the internal space, the former denoting the space-time $E = R^4$ and the latter standing for the set of values of the field, $I = R^n$.

A symmetry transformation can be internal, external or both. Internal symmetry transformations, $\phi^a(x) \rightarrow \phi'^a(x)$, act in the internal space only. Examples are charge conjugation, rotation in flavor or color space of quarks and leptons. External symmetry transformations change the space-time coordinates only, $\phi^a(x) \rightarrow \phi'^a(x')$, eg. the Poincaré group, consisting of space-time translation and Lorentz transformations.

Continuous symmetries generate currents in classical field theories which satisfy the continuity equation according to Noether's theorem. The conserved quantity, the space integral of the time component of the Noether current is usually called charge in case of internal symmetry. The conserved quantities of external symmetries define energy-momentum (translation), angular momentum (space rotation) and a further vector (Lorentz boosts). Note that these symmetry transformations are global, meaning that they are characterized by the same parameters everywhere in the space-time.

B. Local symmetries

It has already been realized in the twenties by H. Weyl and been employed in constructing new theories by Yang and Mills in the sixties that the global symmetries of physics are in conflict with the spirit of special relativity. Let us consider for example a global internal symmetry, represented by the transformation

$$\psi(x) \rightarrow \psi^\star(x) = \omega \psi(x) \quad (15)$$

acting on the multi-component field variable $\phi(x)$ with ω being an element of the symmetry group, typically $\omega \in G = O(n)$ (real fields) or $\omega \in G = \mathfrak{U}(n)$ (complex fields). The symmetry is

global because we have to apply the same change of basis in the internal space anywhere and anytime in the Universe in order to keep the dynamics unchanged. Is the rigid application of the symmetry transformation really necessary in our world where special relativity holds? One can not exchange informations between two locations in the space-time separated by space-like interval, $c^2\Delta t^2 - \Delta \mathbf{x}^2 < 0$ according to special relativity. How can then be a problem in using different bases in the description of physics at space-like separated regions? The symmetry transformations which seem to be in harmony with special relativity should concern change of basis in locations equipped with the possibility of the exchange of physical signals.

The suggestion is to give up any correlations among bases used at different space-time locations and to use local, so called gauge symmetries,

$$\psi(x) \rightarrow \psi^{**}(x) = \omega(x)\psi(x). \quad (16)$$

This is an extreme possibility, opposite to the global transformations. It creates obvious problems if applied to space-time regions with time-like separation, $\Delta t^2 - \Delta \mathbf{x}^2 > 0$ which can exchange signals. In particular, gauge symmetry makes it impossible to obtain any equation of motion for gauge non-invariant quantity which can freely be changed in an arbitrary time-dependent manner. This problem leads to the issue of gauge-fixing not pursued in this simple treatment.

The transformation rule (16) is homogeneous and has the virtue that any local equation

$$0 = F(\psi(x), \chi(x), \dots) \quad (17)$$

which transforms in a homogeneous manner as in (16),

$$F^{**}(\psi(x), \chi(x), \dots) = F(\psi^{**}(x), \chi^{**}(x), \dots) = f(\omega(x))F(\psi(x), \chi(x), \dots), \quad (18)$$

$f(\omega) \neq 0$, remains valid after any gauge transformation. These are called covariant or 'absolute' equations because they are valid in any convention, in other words in any gauge. We are interested in laws of Physics in as simple form as possible. Thus we seek absolute equations. Invariant quantities are called scalars and set of numbers or fields transforming in a homogeneous manner are usually called vectors or tensors. The rules of generating absolute equations consist of prescriptions of constructing scalars, vectors or tensors from scalars, vectors or tensors.

Let us, for the sake of example, consider an imaginary world consisting of two kind of particles, say particle 1 and 2, which participate in an identical manner in their interactions. The field variable has two components and the theory to start with displays a global symmetry group $G = O(2)$ or $G = \mathfrak{O}(2)$. The definition of the particle 1 or 2, amounts to a choice of a basis in the internal

space. It is a convention used by physicists to construct models and communicate the results of their work.

Physicists at different laboratories may use different definitions, called in general conventions below. Experimental physicists need no basis since measurements are performed without making any reference to internal space. Nevertheless they need conventions as soon as they want to compare their findings with model predictions. In this imaginary world the physical phenomena are the same, independently of any choice of conventions. The main question for us in this section is to find the rules of modification of the theory in order to upgrade the global symmetry G into a local one. The result of this procedure, called gauging, is a theory with a gigantic symmetry group, $\mathcal{G} = \otimes \prod_x G_x$. We shall see that the price of such an enlargement of the symmetry is the introduction of a vector field, the gauge field.

Let us start with a theory defined by the Lagrangian $L(\phi, \partial\phi)$, cf. Appendix A, with global symmetry, $\omega \in G$. The Lagrangian has ultra-local terms, involving the field variable $\phi(x)$ at strictly the same space-time point, such as the mass term $\frac{1}{2}m^2\phi_a(x)\phi_a(x)$ or a local potential $\mathfrak{V}(\phi_a(x)\phi_a(x))$. There is no difference between global and local symmetry transformations as far as these terms are concerned. But pieces of the Lagrangian involving space-time derivative of the field are actually detecting the variation of the field on the space-time and are not strictly local. What is important from the point of view of the symmetry is that the transformation rule

$$\partial_\mu\phi(x) \rightarrow \partial_\mu\phi^\bullet(x) = \partial_\mu\omega\phi(x) = \omega\partial_\mu\phi(x) \quad (19)$$

of the global symmetry transformation is modified for local symmetry briefly gauge transformations,

$$\partial_\mu\phi(x) \rightarrow \partial_\mu\phi^\bullet(x) = \partial_\mu\omega(x)\phi(x) = \omega(x)\partial_\mu\phi(x) + (\partial_\mu\omega(x))\phi(x), \quad (20)$$

the trouble maker being the last term. It arises because the derivative compares the field values at neighboring points,

$$\partial_\mu\phi(x) = \lim_{\epsilon \rightarrow 0} \frac{\phi(x + \epsilon n_\mu) - \phi(x)}{\epsilon} \quad (21)$$

and this term represents the contribution due to the different conventions in different points. This contribution should not be there if by difference of the field variables we mean "physical" difference. We should transform the field variables into the same convention before subtraction. The expressing of the field at y into the convention of x , $\phi(y) \rightarrow \omega(x \leftarrow y)\phi(y)$, is a change of basis again. We are interested in this transformation for space-time points within each others vicinity when, the continuous dependence on the space-time coordinate assumed, this transformation is close to the

identity. The possible moves of y into a neighboring x are characterized by an infinitesimal vector $\Delta x^\mu = x^\mu - y^\mu$ and the corresponding change of base

$$\omega(y \leftarrow x) = \mathbb{1} - \Delta x^\mu A_\mu(x) + \mathcal{O}(\Delta^2 x) \approx e^{-\Delta x \cdot A(x)} \quad (22)$$

is given in terms of four generators of the gauge group, $A_\mu(x)$ corresponding to the possible linearly independent moves of the point x . The use of a basis τ^a , $a = 1, \dots, N$ for the Lie-algebra (generators) of an N -dimensional gauge group allows us to write

$$A_\mu(x) = A_\mu^a(x) \tau^a \quad (23)$$

where the basis matrices are assumed to satisfy the normalization conditions

$$\text{tr } \tau^a \tau^b = -\frac{1}{2} \delta^{ab} \quad (24)$$

and commutation relations

$$[\tau^a, \tau^b] = f^{abc} \tau^c. \quad (25)$$

We have a vector field, a gauge field for each direction of the symmetry group following the change of basis one has to compensate for.

The measurable quantities are obviously independent of the choice of basis for the field variable therefore they must be gauge invariant.

C. Covariant derivative

Once we have an expression for the compensation needed to bring the field around a space-time point into the convention at the same point we can define the covariant derivative

$$D_\mu = \partial_\mu + A_\mu \quad (26)$$

as the derivative of the field $\phi(x)$ computed always in the convention at x by

$$\begin{aligned} D_\mu \phi(x) &= \lim_{\epsilon \rightarrow 0} \frac{e^{\epsilon n \cdot A(x + \epsilon n)} \phi(x + \epsilon n_\mu) - \phi(x)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{[1 + \epsilon n \cdot A(x + \epsilon n)] \phi(x + \epsilon n_\mu) - \phi(x)}{\epsilon} \\ &= (\partial_\mu + A_\mu) \phi(x). \end{aligned} \quad (27)$$

The gauge field which appears in the definition of the covariant derivative is sometime called compensating field since its role is to compensate out the contributions of the inhomogeneous

conventions from the derivative of a physical field. It is a generalization of the Lie-derivative, used in the mathematical literature.

Let us now find out the transformation rule for the gauge field $A_\mu(x)$ during the gauge transformation

$$\psi(x) \rightarrow \psi^{**}(x) = \omega(x)\psi(x). \quad (28)$$

The covariant derivative is the derivative of the field computed in fixed convention therefore $D_\mu\phi(x)$ transforms in the same way,

$$D_\mu\psi = (\partial_\mu + A_\mu)\psi \rightarrow D_\mu^{**}\psi^{**} = (\partial_\mu + A_\mu^{**})\psi^{**} = \omega D_\mu\psi = \omega(\partial_\mu + A_\mu)\psi, \quad (29)$$

yielding

$$\omega(\partial_\mu + A_\mu)\psi = (\partial_\mu + A_\mu^{**})\psi^{**} = (\partial_\mu\omega)\psi + \omega\partial_\mu\psi + A_\mu^{**}\omega\psi \quad (30)$$

and

$$A_\mu^{**} = -\partial_\mu\omega\omega^{-1} + \omega A_\mu\omega^{-1}. \quad (31)$$

Let us use the space-time derivative of the identity $\omega(x)\omega^{-1}(x) = \mathbb{1}$,

$$0 = (\partial_\mu\omega)\omega^{-1} + \omega\partial_\mu\omega^{-1}, \quad (32)$$

to write

$$A_\mu \rightarrow A_\mu^{**} = \omega(\partial_\mu + A_\mu)\omega^{-1}. \quad (33)$$

The transformation rule (29) gives the rule of replacing the partial derivative with covariant derivative in the Lagrangian,

$$L(\phi, \partial\phi) \rightarrow L(\phi, D\phi) = L(\phi, (\partial + A)\phi), \quad (34)$$

as the rule of gauging. The interaction induced in this manner between the particle described by the field ϕ and the gauge field is called minimal coupling. It should be clear by inspecting again the derivation of the conserved Noether-current in Chapter A 2 where the new coordinates, related to the global symmetry transformations are actually gauge transformation parameters that the minimal coupling involves the scalar product of the Noether-current and the gauge field, $A_\mu j^\mu$.

D. Parallel transport

Since two internal space vector residing at two different space-time locations can not be compared in their natural bases we need a definition what physically equivalent internal space vectors mean at different space-time points. This is achieved by the parallel transport, a generalization of the construction of the covariant derivative.

Let us consider a continuously derivable path $\gamma^\mu : [0, 1] \rightarrow \mathbb{R}^4$ in the space-time with $\gamma^\mu(0) = x_i^\mu$ and $\gamma^\mu(1) = x_f^\mu$ as initial and final points, respectively and a field $\phi(x)$ defined on this path. What is the condition that the values of this field long our path, $\phi(\gamma(s))$ are physically equivalent, despite the possible dependence of the components of $\phi(\gamma(s))$ on s when expressed in terms of the local basis? Suppose that we have a physical method to check the equivalence of the field along the path. The resulting field $\phi(\gamma(s))$ of a parallel transported internal space vector satisfies the equation

$$\phi(y) = W_\gamma(y, x)\phi(x) \quad (35)$$

where $W_\gamma(y, x)$ is a symmetry (basis) transformation which naturally depends on the choice of the points x and y and a somehow surprising manner will also depend on the path γ , too.

Since this function compensate the change of conventions along the path it should closely be related to the compensating field $A_\mu(x)$ introduced in defining the covariant derivative. In fact, parallel transport of a point of the internal space, vector in short, means that the "physical" components do not change along the path,

$$\frac{d\gamma^\mu}{ds} D_\mu \phi(\gamma(s)) = 0. \quad (36)$$

which can be written as an equation for the parallel transport transformation

$$\frac{d\gamma^\mu}{d\tau} D_{y^\mu} W_\gamma(y, x) = 0 \quad (37)$$

according to Eq. (35). The general solution of this equation is presented in Appendix B, we are now satisfied by the approximate solution for short paths,

$$W_\gamma(x + \Delta x, x) = e^{-\Delta x^\mu A_\mu(x)} \quad (38)$$

which naturally agrees with Eq. (B1) in the limit $x - y \rightarrow 0$.

It is easy to find out the transformation rule for parallel transport under gauge transformations. The starting point is that the product $\phi^\dagger(y)W_\gamma(y, x)\phi(x)$ or $\phi(y)W_\gamma(y, x)\phi(x)$ for complex or real ϕ , respectively, is gauge invariant. Hence the equation $\omega^{-1}(y)W_\gamma^\star(y, x)\omega(x) = W_\gamma(y, x)$ follows

and we have

$$W_{\gamma}^{\bullet\bullet}(y, x) = \omega(y)W_{\gamma}(y, x)\omega^{-1}(x). \quad (39)$$

An unusual feature of the parallel transport $W_{\gamma}(y, x)$ is its path dependence. To understand the impact of this dependence let us imagine first a gauge field $A_{\mu}(x)$ for which $W_{\gamma}(y, x)$ is independent of the path γ . In this case we may choose a reference point in space-time, say x_0 and extend its convention, basis to the rest of the space-time by performing the gauge transformation $\omega(x) = W^{-1}(x, x_0)$. This transformation renders all parallel transport trivial since

$$W(y, x) = W^{-1}(y, x_0)W(y, x)W(x, x_0) = W(x_0, y)W(y, x_0) = \mathbb{1} \quad (40)$$

in this gauge. The gauge field which leads such a path independent parallel transports is called pure gauge because it can be canceled by an appropriate gauge transformation.

There is an equivalent characterization of path independence of parallel transports, it is the triviality of parallel transport on closed paths, $W_{\gamma}(x, x) = \mathbb{1}$. In fact, let us choose another point than x of the closed path γ what will be denoted by $y = \gamma(s)$, $0 < s < 1$. We introduce the fragments of γ from x to y and from y to x , as $\gamma_1(t) = \gamma(ts)$ and $\gamma_2(t) = \gamma(s + t(1 - s))$ which satisfy the equation

$$W_{\gamma}(x, x) = W_{\gamma_1}(x, y)W_{\gamma_2}(y, x) = W_{\gamma_1}(x, y)W_{\gamma_2}^{-1}(x, y) \quad (41)$$

the two parallel transports, $W_{\gamma_1}(x, y)$ and $W_{\gamma_2}^{-1}(y, x)$ correspond to two different paths connecting the events x and y therefore the path dependence is equivalent with the non-triviality of the parallel transports along closed paths.

E. Field strength tensor

The gauging, the upgrade of a global symmetry to a local one brings in a generator valued vector field. We are accustomed to the fact that fields corresponds to particles. Therefore the gauging of a symmetry suggests the presence of spin 1 bosons in the system. The dynamics of these particle can not come from the Lagrangian (34) because of the lack of the velocities $\partial_0 A_{\mu}$ in it. The simplest solution is the add a new term to the Lagrangian $L \rightarrow L + L_A$ where L_A satisfies the following conditions:

1. It should be **quadratic in the velocities**, $L_A = \mathcal{O}((\partial_0 A_{\mu})^2)$.
2. It should be **Lorentz invariant**.

3. It should be **gauge invariant**.

The last property requires that L_A should be vanishing for pure gauge fields, it should depend on the non-pure gauge component of the field. This property suggests L_a , a local quantity, be constructed in terms of the deviation of parallel transports on infinitesimally small loops from the identity. Let us therefore consider the parallel transport of a field along a rectangle $x \rightarrow x + u \rightarrow x + u + v \rightarrow x + v \rightarrow x$ in space-time, u and v being infinitesimal, non-parallel vectors. The change of the field during the parallel transport is infinitesimal, as well, $\phi \rightarrow \phi + \delta\phi$ and should be linear in u , v and ϕ itself. Therefore one expects the relation

$$\delta\phi^a = -F_{b\mu\nu}^a u^\mu v^\nu \phi^b. \quad (42)$$

According to the definition of the parallel transport along an infinitesimal straight line, Eq. (38), the parallel transport along the rectangle is

$$\mathcal{P}_\square = e^{vA(x)} e^{uA(x+v)} e^{-vA(x+u)} e^{-uA(x)}. \quad (43)$$

We expand up to the displacement squares for each exponential functions gives

$$\begin{aligned} \mathcal{P}_\square \approx & \left(\mathbb{1} + vA(x) + \frac{1}{2}[vA(x)]^2 \right) \left(\mathbb{1} + uA(x+v) + \frac{1}{2}[uA(x+v)]^2 \right) \\ & \times \left(\mathbb{1} - vA(x+u) + \frac{1}{2}[vA(x+u)]^2 \right) \left(\mathbb{1} - uA(x) + \frac{1}{2}[uA(x)]^2 \right) \end{aligned} \quad (44)$$

which can be simplified within this approximation to

$$\begin{aligned} \mathcal{P}_\square \approx & \mathbb{1} + (v\partial)uA - (u\partial)vA - (uA)(vA) + (vA)(uA) \\ = & \mathbb{1} - u^\mu v^\nu (\partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]), \end{aligned} \quad (45)$$

resulting the field strength tensor

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu] = [D_\mu, D_\nu], \quad (46)$$

a generator valued field,

$$F_{\mu\nu} = F_{\mu\nu}^a \frac{\tau^a}{2i}, \quad (47)$$

with

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c. \quad (48)$$

The gauge field strength is the measure of the non-triviality of the parallel transport of a state along a closed path.

The transformation rule for the parallel transport on a closed path,

$$\mathbb{1} - u^\mu v^\nu F_{\mu\nu}(x) \rightarrow \omega(x)[\mathbb{1} - u^\mu v^\nu F_{\mu\nu}(x)]\omega^{-1}(x) \quad (49)$$

gives the transformation rule

$$F_{\mu\nu}(x) \rightarrow \omega(x)F_{\mu\nu}(x)\omega^{-1}(x). \quad (50)$$

One can show that the triviality of the parallel transport over infinitesimal loops, the vanishing of the field strength tensor, assures the triviality of the parallel transport over arbitrary loops and restricts the gauge field to a pure gauge form, $A_\mu = \omega\partial_\mu\omega^{-1}$ in space-time without boundary conditions. When boundary conditions apply then the elimination of a pure gauge field configuration may be impossible. Such a configuration play an important role in the dynamics.

The unique solution of the constraints for L_A on a space-time with trivial topology is the Yang-Mills Lagrangian,

$$L_{YM} = \frac{1}{2g^2}\text{tr}(F_{\mu\nu})^2 = -\frac{1}{4g^2}(F_{\mu\nu}^a)^2, \quad (51)$$

which is fixed up to the coupling constant g . It is advantageous to use the notation $A_\mu \rightarrow gA_\mu$ in perturbation expansion, giving

$$L_{YM} = -\frac{1}{4}F_{\mu\nu}^a F^{\mu\nu a}. \quad (52)$$

III. GRAVITY

It seems to be an essential feature of Nature that all known interactions belong to the class of gauge theories. For the electromagnetic, weak and strong interactions the internal space is independent of the space-time. The special feature of gravity is that it influences the geometry of the space-time therefore its internal space is not independent of its external space. The formalism of general relativity will be introduced below by underlying its origin in gauge theories.

A. Classical field theory on curved space-time

The mathematical view of a field configurations, $\phi(x)$, is a map $\phi : E \rightarrow I$ which describes “what” (I) happens “where” (E). Gravity and other interactions provide dynamics for E and I , respectively. Such a dynamics may generate singular x -dependence which can be avoided by renouncing the global, single valued nature of the dynamical fields. Instead, the fields are expected

to produce a well defined image point in subsets of the space-time only and the resulting structure is called differentiable manifold. The regions $\mathcal{U}_j \subset E$ where both the space-time coordinates and the fields have unique, well defined values are maps and the collection of maps, $\{\mathcal{U}_j\}$, an atlas, is supposed to satisfy the following properties:

1. **Maps:** Each space-time point correspond at least to one map. ie. each space-time point can be identified by means of the coordinates.
2. **Coordinates:** There is a one-to-one mapping, $x_j : \mathcal{U}_j \rightarrow V$, of each map into an open subset V of R^{d_E} . These d_E -dimensional functions, $x_j^\mu(p)$, $\mu = 1, \dots, d_E$, $p \in \mathcal{M}$, play the role of coordinates defined for each map and the space-time looks locally d_E -dimensional, in agreement with the Equivalence Principle. We assume $d_E = 4$ in what follows. Experimental devices are supposed to be available to measure the values of the coordinates at each space-time point.
3. **Coordinate transformations:** The coordinates obtained in the overlapping region of two maps depend on each other on an infinitely many times differentiable manner and the domain of definition of these functions is always an open set in R^4 .

To emphasize the local nature of interactions one associates an internal space, I_p , to each space-time point $p \in E$. The dimension of this space may be rather large, it is the sum of the number of components of all dynamical fields of our theory. It is advantageous separate two components of the internal space, $I = T \oplus F$. T , the tangent space is d_E -dimensional, it consists of the possible tangent vectors of world lines at the given space-time and is reserved for the description of the space-time geometry. The remaining $d_I - d_E$ dimensions of F correspond to the non-gravitational fields, for instance $d_I - d_E = 4$ for classical electrodynamics.

One may give a formal definition of T_p , it is the collection of equivalence classes of world lines $x(s)$, crossing p , $x(s_p) = x(p)$ where $x(s)$ and $x'(s)$ are considered equivalent if they have the same tangent vector in p , $\dot{x}(s_p) = \dot{x}'(s_p)$. We assume that experimental devices are available at each space-time point to measure the four velocity of point particles and identify a point in the tangent space. The gravitational interaction determines the motion masses, therefore it governs the dynamics in T .

An important result of this particular construction of the tangent space is that the vectors $u \in T_p$, can be related to the infinitesimal displacement, $p \rightarrow p'$,

$$x_j^\mu(p') = x_j^\mu(p) + \delta s u^\mu, \quad (53)$$

where $u^\mu = \dot{x}^\mu(s)$ is the tangent vector of a curve, traversing the point p . This equation reflects the relation between the tangent space and tangent vectors but remains formal. Both the calculation of the tangent vectors and the construction of coordinate systems become well defined by choosing a standard map, $X(p)$, for the neighborhood of p and using eq. (53) to define the map $x(p') = x(p) + \delta x$ by the equation

$$X(p') = X(p) + \delta x^\mu e_\mu(p), \quad (54)$$

involving the coordinate basis vectors,

$$e_\mu(p) = \partial_\mu X(x), \quad (55)$$

using the coordinate transformation $X(x)$.

The change of the coordinate system is a local transformation of the vicinity of the point p . One can treat such a local change of our conventions in the labeling of the points as a gauge transformation. This is the point where gauge symmetry makes its appearance in differential geometry. The basis vectors of another map, $x'(p')$, are related to e_μ by the equation

$$e_\mu = e'_\nu \frac{\partial x'^\nu}{\partial x^\mu}. \quad (56)$$

The corresponding transformation rule of a contravariant vector is

$$e^\mu = \frac{\partial x^\mu}{\partial x'^\nu} e'^\nu \quad (57)$$

These equations are based on a particular standard map. Its change, $X(p) \rightarrow X'(p)$, generates a linear transformation of the component of the basis vectors, without touching their covariant index,

$$e'_\mu = \frac{\partial X'}{\partial X^\nu} (e_\mu)^\nu. \quad (58)$$

The transformation rules, (56)-(57), can easily be memorized by the following observations: One could have derived eq. (56) by using (55) by dropping $X(x)$. The result is that the covariant vectors transform as partial derivatives,

$$\partial_\mu = \partial'_\nu \frac{\partial x'^\nu}{\partial x^\mu}. \quad (59)$$

Since $dx^\mu \partial_\mu$ remains invariant under coordinate transformations the contravariant vectors transform as the infinitesimal coordinate changes,

$$\delta x^\mu = \frac{\partial x^\mu}{\partial x'^\nu} \delta x'^\nu. \quad (60)$$

This observation is the starting point of an elegant formalism in differential geometry. It sometime is called the absolute calculus since these objects can formally be defined without using a standard (local) coordinate system. Equations which preserve their form under the reparametrization of the coordinates are called covariant. The absolute calculus produces covariant equations, constructed in terms of functions (coordinates), their derivatives (covariant vector fields) and infinitesimal changes (contravariant vector fields). One arrives at nice and compact equations in this manner, showing the geometrical essence in as clear manner as possible. But this scheme has two drawbacks from the point of view of physics. One is that one is interested in the agreement between observed and theoretically predicted quantities and such a comparison can not be made without actually “dirtying our hand” with a given coordinate system. The other is that while the idea of representing vector fields by partial derivatives or by infinitesimal changes is correct and justified mathematically it leads to a wrong intuition in physics. I believe that it is more advantageous first to spend the time needed by improving our way of reading and understanding equations, written in terms of a coordinate system until we recognize the general structure. When the physics is properly expressed then, as a second step, one may go over the coordinate independent scheme if one wishes.

The tangent space represents vectors at a given space-time point which may appear in physical laws. One introduces different tangent spaces at different space-time point because their elements, the vectors, are defined for each space-time point in an independent manner, locally, by physical measurements. When this construction is restricted to gravitational interaction only, $d_I = d_E$ and $F = 0$, it gives a differentiable manifold, the mathematical framework of General Relativity.

B. Geometry

There are three important properties of the space-time geometry which appears in gravitational interactions.

1. The **metric** property is related to the existence of an invariant distance and it is essential in establishing spatial and temporal distances between events. It is assumed that physical measurements provide us spatial distances, time intervals and angles to determine the metric. Due to the Equivalence Principle the metric structure must locally be compatible with Special Relativity. This is achieved by introducing the invariant length in a local manner,

$$ds^2(x) = dx^\mu g_{\mu\nu}(x) dx^\nu \quad (61)$$

where the metric tensor $g_{\mu\nu}(x)$ is a symmetric tensor with three positive and a negative

eigenvalues, i.e. with signature $+, -, -, -$.

2. The **affine** property controls parallelism by determining what directions can we consider parallel at different space-time points. The velocity and the internal angular momentum, the classical spin, of a small enough body is constant in the absence of external forces and the acceleration and the time derivative of the spin are vanishing. According to the Equivalence Principle this remains valid locally in a suitable chosen coordinate system in the presence of external gravitational field. Thus the velocity and the spin preserve their directions in a non-trivial geometry and can be used to establish the concept of parallelism. The affine structure is realized by the affine connection, alias compensating or gauge field of gauge theories.
3. The **torsion** of the space-time represents certain distortions of the space-time. This property seems to be important in describing the interaction of the quantum mechanical spin with the gravitational field only. Note that though the Equivalence Principle has been build in by point 2. in Section III A, it can eventually be violated by implying the torsion in the dynamics because the distortions, characterized by the torsion can not be eliminated by using suitable local coordinate system. We consider General Relativity in classical physics and the torsion will be assumed to be absent.

C. Gauge group

It is instructive to recall that the absolute location in space-time becomes unobservable and the space-time location is relative when the dynamics is translation invariant. The spatial directions are relative in rotational invariant systems. The symmetry with respect to Lorentz boost makes the absolute velocity unobservable in Special Relativity. The acceleration and all higher order derivatives of the world line are rendered relative in General Relativity by imposing invariance of the dynamics under reparametrization of the space-time coordinates, called space-time diffeomorphism. Hence the choice of the coordinate system in the space-time is mere convention, the role of the coordinates is to identify space-time points only and the actual numerical values of the coordinate have no physical meaning.

The fundamental physical laws. eg. the Maxwell-equations, are supposed to be local in space-time, they can be expressed as equations among dynamical quantities, corresponding to the same space-time location. There is indeed no need of absolute coordinates for such a description. When

we do not intend to follow and find the equation of motion for all dynamical degree of freedom then we seek an effective description. The motion of point particles or propagation of waves which are not followed by us relate and correlate dynamical quantities at different space-time location and are used to introduce direction and distance in space-time. We are interested here the fundamental laws therefore we assume that the equations of motions are expressed for each space-time location independently. This assumption leads to a rich gauge theory structure in an obvious manner and gravity can be founded as a gauge theory with either external or internal symmetries.

1. Space-time diffeomorphism

The reparametrization of the coordinates, the diffeomorphism of the external space, can be generated by the infinitesimal local translations in space time, $x^\mu(x) \rightarrow x'^\mu(x) = x^\mu(x) + \delta x^\mu(x)$. If the value of the coordinates are not relevant then the change of the coordinate system is represented by the change of the tangent vectors of world lines, in particular the coordinate axes. Therefore, the gauge group is $GL(4)$, consisting of 4×4 non-singular matrices describing the transformation of the coordinate axes $e_\mu(x)$ during general coordinate transformations,

$$e_\mu(x) \rightarrow e'_\mu(x) = M_\mu{}^\nu(x)e_\nu(x). \quad (62)$$

The affine structure of the space-time which is the central feature of this formalism handles the relation of directions at different space-time points by means of parallel transport in the $GL(4)$ gauge theory. The drawback of this line of thought is that the other independent geometrical structure, the metric which is a key player in the traditional approach to General Relativity is constructed in an indirect manner from the affine connection.

The internal space is chosen to be the tangent space, the directional vectors at each point of the space-time and its elements are contravariant vectors. The independent field variables are the metric tensor $g_{\mu\nu}$ and the $GL(4)$ gauge field, $\Gamma^\rho{}_{\mu\nu}$, the affine connection. The metric tensor is symmetrical, contains 10 independent component and has the signature $(+, -, -, -)$. Once the metric tensor is introduced the tangent spaces can be represented covariant vectors, too. Furthermore, the direct product of the tangent space gives rise to the space of local tensors, as well.

The representation of the diffeomorphism by the transformation of the tangent spaces, (62) with $M_\mu{}^\nu(x) = \delta_\mu^\nu - \partial_\mu \delta x^\nu$ raises a consistency issue, namely what conditions should the four vector fields, $e_\mu(x)$ satisfy to make up a coordinate basis? The necessary condition arises from the

symmetry of the second partial derivatives, $\partial_\mu e_\nu = \partial_\mu \partial_\nu x' = \partial_\nu \partial_\mu x' = \partial_\nu e_\mu$. It is easy to check that this condition is sufficient. In fact,

$$\partial_\mu e_\nu = \partial_\nu e_\mu \quad (63)$$

is sufficient to assure the local existence and unicity of integral curves, the solution of the equations $\partial_\mu x'(x) = e_\mu(x)$. The tetrads, satisfying (63) and can be used to construct well defined local coordinates are called holonomic. The possibility of representing the reparametrization of the space-time by the transformation of the coordinate basis vectors stems from the preservation of the holonomy under diffeomorphism.

2. Internal Poincaré group

Both the metric and the affine structures can be derived in the gauge theory formalism by means of internal gauge symmetry. The starting point is the Equivalence Principle which assures the existence of a coordinate system with local Lorentz invariance.

Lorentz transformations: The local Lorentz coordinate axes, $e^a(x)$, can be considered as basis in the internal Lorentz space. But one might as well use another reference frame, obtained by a local Lorentz transformation,

$$e^a(x) \rightarrow e'^a(x) = \omega^a_b(x) e^b(x). \quad (64)$$

Thus one is led to propose Lorentz transformation as a local symmetry. A gauge field, $\omega^a_{b\mu}$, introduced to handle the compensations of the local Lorentz transformations defines the affine structure of the space-time.

Translations: The special feature of gravity is the relation between dynamics and geometry, a link between the internal and the external spaces. The external diffeomorphism, the coordinate reparametrization invariance can be generated by the infinitesimal local translations in space time, $x^\mu(x) \rightarrow x'^\mu(x) = x^\mu(x) + \delta x^\mu(x)$. The tangent space, T_p , consists of tangent vectors of world lines, \dot{x}^μ , at p . By representing the tangent vectors in the local Lorentz reference frame, $\dot{x}^\mu = \dot{\xi}^a e_a^\mu$, we turn this latter into the tangent space, T_p . The internal space equivalent of local, infinitesimal translations,

$$\xi^a(x) \rightarrow \xi'^a(x) = \xi^a(x) + \delta \xi^a(x) \quad (65)$$

is now considered as a local gauge transformation. The gauge field compensating such a local modification of the coordinates of the local Lorentz spaces, $e_\mu^a(x)$ is called vierbein and provides

the desired link between infinitesimal shifts in the internal Lorentz space and the space-time:

$$\delta\xi^a(x) = e_\mu^a \delta x^\mu(x). \quad (66)$$

We shall use Greek and Latin letters to denote vector indices in the Lorentz and in the space-time coordinate system, respectively. The translations (65), together with the Lorentz transformations (64) form the Poincaré group as gauge symmetry.

An unexpected bonus of the Poincaré group formalism is the possibility of treating fermions. In fact, fermions do not have well covariant transformations properties under the $GL(4)$ group, the general change of coordinates. They show well defined transformation rules for the Lorentz group only. Another advantageous feature of this formalism is the natural way torsion couples to angular momentum in the dynamics.

D. Gauge theory of diffeomorphism

The simpler framework for gravity, based on the external diffeomorphism as gauge group is introduced first.

1. Covariant derivative

The affine structure is defined by the connection Γ_μ , a 4×4 matrix valued vector field with 64 independent components. We shall use the notation $(\Gamma_\rho)^\mu{}_\nu = \Gamma^\mu{}_{\nu\rho}$ for the components of the connection. The covariant derivative,

$$D_\nu v^\mu = \partial_\nu v^\mu + \Gamma^\mu{}_{\rho\nu} v^\rho, \quad (67)$$

detects the 'real', physical changes of a vector field by projecting out the changes of the vector components arising from the changing conventions. By suppressing the indices in Eq. (67) we have for contravariant vectors

$$(D_\nu v)^\mu = (\partial_\nu v + \Gamma_\nu v)^\mu. \quad (68)$$

The metric structure is behind the reduplication of the vectors and suggests the definition

$$(D_\nu v)_\mu = (\partial_\nu v - v\Gamma_\nu)_\mu \quad (69)$$

for covariant vector field. The reason is that this definition allows us the contraction of indices within the covariant derivative and to recover the equivalence of the covariant and the partial

derivatives for scalar fields,

$$D_\mu(u_\nu v^\nu) = (\partial_\mu u - u\Gamma_\mu)_\nu v^\nu + u_\nu(\partial_\mu v + \Gamma_\mu v)^\nu = \partial_\mu(u_\nu v^\nu) \quad (70)$$

The action of the covariant derivative is extended over any tensor field by performing the necessary compensation on each vector index, eg.

$$D_\nu v_\rho^\mu = \partial_\nu v_\rho^\mu + \Gamma_{\kappa\nu}^\mu v_\rho^\kappa - v_\kappa^\mu \Gamma_{\rho\nu}^\kappa. \quad (71)$$

It is easy to check that such an extension reproduces Leibniz rule,

$$D_\mu(u^\rho v^\sigma) = (D_\mu u^\rho)v^\sigma + u^\rho D_\mu(v^\sigma). \quad (72)$$

Notice that a coordinate transformation $x \rightarrow x'(x)$ induces the change

$$\Gamma_{\nu\rho}^\mu \rightarrow \Gamma'_{\nu\rho}{}^\mu = \frac{\partial x^\sigma}{\partial x'^\rho} \omega_\kappa^\mu (\delta_\lambda^\kappa \partial_\sigma + \Gamma_{\lambda\sigma}^\kappa) (\omega^{-1})^\lambda{}_\nu. \quad (73)$$

according to the transformation rule (33) and the application of the second equation in (56) to the connection as a four-vector. The expressions $\omega_\kappa^\mu = \frac{\partial x'^\mu}{\partial x^\kappa}$ and $(\omega^{-1})^\mu{}_\kappa = \frac{\partial x^\mu}{\partial x'^\kappa}$ allow us to write

$$\begin{aligned} \Gamma_{\nu\rho}^\mu &\rightarrow \frac{\partial x^\sigma}{\partial x'^\rho} \frac{\partial x'^\mu}{\partial x^\kappa} \frac{\partial^2 x^\kappa}{\partial x'^\nu \partial x^\sigma} + \frac{\partial x^\sigma}{\partial x'^\rho} \frac{\partial x'^\mu}{\partial x^\kappa} \Gamma_{\lambda\sigma}^\kappa \frac{\partial x^\lambda}{\partial x'^\nu} \\ &= \frac{\partial x'^\mu}{\partial x^\kappa} \frac{\partial^2 x^\kappa}{\partial x'^\nu \partial x'^\rho} + \frac{\partial x^\sigma}{\partial x'^\rho} \frac{\partial x'^\mu}{\partial x^\kappa} \Gamma_{\lambda\sigma}^\kappa \frac{\partial x^\lambda}{\partial x'^\nu}, \end{aligned} \quad (74)$$

showing that the affine connection is not a tensor due to the inhomogeneous term in the gauge transformation (33), the first term on the right hand side of (74). But the antisymmetric part in the covariant indices, called torsion,

$$S_{\nu\mu}^\rho = \frac{1}{2}(\Gamma_{\mu\nu}^\rho - \Gamma_{\nu\mu}^\rho). \quad (75)$$

is a tensor. The form (31) of the gauge transformation is sometime useful,

$$\begin{aligned} \Gamma_{\nu\rho}^\mu \rightarrow \Gamma'_{\nu\rho}{}^\mu &= -\frac{\partial x^\sigma}{\partial x'^\rho} \partial_\sigma \omega_\kappa^\mu \delta_\lambda^\kappa (\omega^{-1})^\lambda{}_\nu + \frac{\partial x^\sigma}{\partial x'^\rho} \omega_\kappa^\mu \Gamma_{\lambda\sigma}^\kappa (\omega^{-1})^\lambda{}_\nu \\ &= -\frac{\partial x^\kappa}{\partial x'^\nu} \frac{\partial^2 x'^\mu}{\partial x^\kappa \partial x^\sigma} \frac{\partial x^\sigma}{\partial x'^\rho} + \frac{\partial x^\sigma}{\partial x'^\rho} \frac{\partial x'^\mu}{\partial x^\kappa} \Gamma_{\lambda\sigma}^\kappa \frac{\partial x^\lambda}{\partial x'^\nu}. \end{aligned} \quad (76)$$

It is advantageous to use harmonic gauge, defined by

$$\Gamma^\rho = g^{\mu\nu} \Gamma_{\mu\nu}^\rho = 0 \quad (77)$$

for solving the equations of motion. The name comes from the equation

$$\square x^\mu = g^{\rho\nu} D_\rho D_\nu x^\mu = g^{\nu\rho} D_\rho \partial_\nu x^\mu = -\Gamma^\mu \quad (78)$$

where the second equation holds because x^μ is a scalar field for a given value of μ , stating that the coordinates are harmonic functions. Eq. (76) leads to the transformation rule

$$\begin{aligned}\Gamma^\mu \rightarrow g^{\nu\rho}\Gamma_{\nu\rho}^{\prime\mu} &= g^{\tau\sigma}\frac{\partial x^{\prime\nu}}{\partial x^\tau}\frac{\partial x^{\prime\rho}}{\partial x^\sigma}\left(-\frac{\partial x^\kappa}{\partial x^{\prime\nu}}\frac{\partial^2 x^{\prime\mu}}{\partial x^\kappa\partial x^\sigma}\frac{\partial x^\sigma}{\partial x^{\prime\rho}}+\frac{\partial x^\sigma}{\partial x^{\prime\rho}}\frac{\partial x^{\prime\mu}}{\partial x^\kappa}\Gamma_{\lambda\sigma}^\kappa\frac{\partial x^\lambda}{\partial x^{\prime\nu}}\right) \\ &= -g^{\tau\sigma}\frac{\partial^2 x^{\prime\mu}}{\partial x^\tau\partial x^\sigma}+\frac{\partial x^{\prime\mu}}{\partial x^{\prime\kappa}}\Gamma^\kappa\end{aligned}\quad (79)$$

which gives the equation

$$g^{\tau\sigma}\frac{\partial^2 x^{\prime\mu}}{\partial x^\tau\partial x^\sigma}=\frac{\partial x^{\prime\mu}}{\partial x^{\prime\kappa}}\Gamma^\kappa.\quad (80)$$

The harmonic gauge can always be reached by solving this equation for $x^{\prime\mu}(x)$ when the field Γ^κ is given.

The Equivalence Principle can be rephrased in a mathematical form. Let us first consider a gauge theory where the gauge field transforms according to eq. (33). It is easy to see that the gauge transformation, $\omega(x) = e^{(x^\mu - x_0^\mu)A_\mu(x_0)}$, defined by the help of a given, fixed space-time point, x_0 , yields

$$\begin{aligned}A_\mu^{\bullet\bullet}(x) &= \omega(x)(\partial_\mu + A_\mu(x))\omega^{-1}(x) \\ &= [\mathbb{1} + (x^\mu - x_0^\mu)A_\mu(x_0)](\partial_\mu + A_\mu(x))[\mathbb{1} - (x^\mu - x_0^\mu)A_\mu(x_0)] + \mathcal{O}((x - x_0)^2) \\ &= \mathcal{O}(x - x_0).\end{aligned}\quad (81)$$

In other words, the gauge field can be eliminated at x_0 by means of a suitable gauge transformation, rendering the covariant derivative locally equivalent with the partial derivative, and leaving only its space-time derivatives non-vanishing. In case of gravity the internal and external spaces are related and we first perform linear change of coordinates in such a manner that the metric tensor assumes its Minkowski form at x_0 . After that we make a further nonlinear coordinate transformation, $x \rightarrow x'$, given by

$$x^\mu - x_0^\mu = x^{\prime\mu} - x_0^{\prime\mu} - \frac{1}{2}\Gamma_{\nu\rho}^\mu(x_0)(x^{\prime\nu} - x_0^{\prime\nu})(x^{\prime\rho} - x_0^{\prime\rho})\quad (82)$$

where

$$\frac{\partial x^\kappa}{\partial x^{\prime\mu}} = \delta_\mu^\kappa - \frac{1}{2}\Gamma_{\mu\rho}^\kappa(x_0)(x^{\prime\rho} - x_0^{\prime\rho}) - \frac{1}{2}\Gamma_{\nu\mu}^\kappa(x_0)(x^{\prime\nu} - x_0^{\prime\nu}),\quad (83)$$

and in particular

$$\frac{\partial x^\kappa}{\partial x^{\prime\mu}}\Big|_{x'=x_0'} = \delta_\mu^\kappa.\quad (84)$$

The transformation rule of the affine connection,

$$\Gamma'^{\mu}_{\nu\rho}(x') = -\frac{\partial x'^{\mu}}{\partial x^{\kappa}}\Gamma^{\kappa}_{\nu\rho}(x_0) + \frac{\partial x^{\sigma}}{\partial x'^{\rho}}\frac{\partial x'^{\mu}}{\partial x^{\kappa}}\Gamma^{\kappa}_{\lambda\sigma}(x)\frac{\partial x^{\lambda}}{\partial x'^{\nu}}, \quad (85)$$

gives at $x = x_0$

$$\Gamma'^{\mu}_{\nu\rho}(x'_0) = -\Gamma^{\mu}_{\nu\rho}(x_0) + \Gamma^{\mu}_{\nu\rho}(x_0) = 0. \quad (86)$$

In other words, the metric tensor can be brought into its flat space-time form and the affine connection can be made vanishing at any fixed space-time point by the use of appropriate coordinates. Such an elimination of the non-trivial geometry of the space time is a local feature because the second derivatives of the metric tensor and the first derivatives of the affine connection remain non-trivial in any coordinate system.

2. Lie derivative

The Lie derivative, the covariant derivative, generated by space-time diffeomorphism, gives the change of a field $\phi(x)$ during a space-time diffeomorphism $x^{\mu} \rightarrow x^{\mu} - w^{\mu}(x)$ expressed in the coordinate basis at x . The field is symmetric under such a diffeomorphism if its Lie derivative is vanishing.

Let us consider for the sake of simplicity a vector field $u^{\nu}(x)$. Its Lie derivative with respect to the diffeomorphism $w(x)$ is the sum of two terms. The first is the change $u \rightarrow u(x + w)$, induced by the coordinate transformation, $w^{\nu}\partial_{\nu}u^{\mu}$. The other contribution comes from the transformation of the vector $u(x + w)$ into the basis at x . The corresponding transformation (57) contains the matrix $\omega^{\mu}_{\nu} = \delta^{\mu}_{\nu} - \partial_{\nu}w^{\mu}$. The Lie derivative is therefore

$$\nabla_w u^{\mu} = w^{\nu}\partial_{\nu}u^{\mu} - \partial_{\nu}w^{\mu}u^{\nu}. \quad (87)$$

The form

$$\nabla_w u^{\mu} = w^{\nu}D_{\nu}u^{\mu} - u^{\nu}D_{\nu}w^{\mu} + 2S^{\mu}_{\rho\nu}u^{\rho}w^{\nu} \quad (88)$$

shows that the Lie derivative of vector field is a covariant vector field. The Lie derivative of a scalar is given by the partial derivative, $\nabla_w \phi = w^{\nu}\partial_{\nu}\phi$ and the generalization of (88) for covariant vectors and tensors is straightforward, eg.

$$\begin{aligned} \nabla_w u^{\mu}_{\kappa} &= w^{\nu}\partial_{\nu}u^{\mu}_{\kappa} - \partial_{\nu}w^{\mu}u^{\nu}_{\kappa} + \partial_{\kappa}w^{\nu}u^{\mu}_{\nu} \\ &= w^{\nu}D_{\nu}u^{\mu}_{\kappa} - u^{\nu}_{\kappa}D_{\nu}w^{\mu} + u^{\mu}_{\nu}D_{\kappa}w^{\nu} + 2S^{\mu}_{\rho\nu}u^{\rho}_{\kappa}w^{\nu} + 2S^{\rho}_{\kappa\nu}u^{\mu}_{\rho}w^{\nu}. \end{aligned} \quad (89)$$

3. Field strength tensor

The $GL(4)$ field strength tensor is

$$F_{\mu\nu} = [D_\mu, D_\nu] = [\partial_\mu + \Gamma_\mu, \partial_\nu + \Gamma_\nu] = \partial_\mu \Gamma_\nu - \partial_\nu \Gamma_\mu + [\Gamma_\mu, \Gamma_\nu] \quad (90)$$

which is antisymmetric in the space-time indices,

$$F_{\mu\nu} = -F_{\nu\mu}. \quad (91)$$

The field strength tensor is called the curvature tensor and reads

$$R^\mu{}_{\nu\rho\sigma} = (F_{\rho\sigma})^\mu{}_\nu = \partial_\rho \Gamma^\mu{}_{\nu\sigma} - \partial_\sigma \Gamma^\mu{}_{\nu\rho} + \Gamma^\mu{}_{\kappa\rho} \Gamma^\kappa{}_{\nu\sigma} - \Gamma^\mu{}_{\kappa\sigma} \Gamma^\kappa{}_{\nu\rho} \quad (92)$$

with all indices shown. According to the remark, made after Eq. (50) the vanishing of the curvature tensor is equivalent with the absence of gravitational forces in a space-time without boundary.

A useful identity for the curvature tensor obtained from symmetrical connection is

$$R^\rho{}_{\kappa\mu\nu} + R^\rho{}_{\mu\nu\kappa} + R^\rho{}_{\nu\kappa\mu} = 0. \quad (93)$$

An important relation for the curvature tensor follows from the Bianchi identity for commutators,

$$0 = [A, [B, C]] + [B, [C, A]] + [C, [A, B]], \quad (94)$$

which yields

$$\begin{aligned} 0 &= [D_\mu, [D_\nu, D_\rho]] + [D_\nu, [D_\rho, D_\mu]] + [D_\rho, [D_\mu, D_\nu]] \\ &= [D_\mu, F_{\nu\rho}] + [D_\nu, F_{\rho\mu}] + [D_\rho, F_{\mu\nu}] \\ &= D_\mu F_{\nu\rho} + D_\nu F_{\rho\mu} + D_\rho F_{\mu\nu}, \end{aligned} \quad (95)$$

or

$$0 = D_\mu R^\sigma{}_{\kappa\nu\rho} + D_\nu R^\sigma{}_{\kappa\rho\mu} + D_\rho R^\sigma{}_{\kappa\mu\nu}, \quad (96)$$

by writing all indices explicitly.

The Lagrangian of a gauge field is usually a quadratic expression of the the field strength tensor, $\text{tr} F_{\mu\nu} F^{\mu\nu}$. The distinguished feature of gravity is that its internal space, the tangent space of external space, is related to the space-time. This feature allows us to contract internal index

with external one and to construct invariant expressions which are linear in the field strength. We may make three different contractions, the Ricci tensors, defined as

$$\begin{aligned} R_{\nu\sigma} &= R^{\rho}{}_{\nu\rho\sigma} \\ &= \partial_{\rho}\Gamma^{\rho}{}_{\nu\sigma} - \partial_{\sigma}\Gamma^{\rho}{}_{\nu\rho} + \Gamma^{\rho}{}_{\kappa\rho}\Gamma^{\kappa}{}_{\nu\sigma} - \Gamma^{\rho}{}_{\kappa\sigma}\Gamma^{\kappa}{}_{\nu\rho} \end{aligned} \quad (97)$$

$R^{\rho}{}_{\nu\sigma\rho} = -R_{\nu\sigma}$, and

$$\begin{aligned} R'_{\rho\sigma} &= R^{\mu}{}_{\mu\rho\sigma} \\ &= \partial_{\rho}\Gamma^{\mu}{}_{\mu\sigma} - \partial_{\sigma}\Gamma^{\mu}{}_{\mu\rho} + \Gamma^{\mu}{}_{\kappa\rho}\Gamma^{\kappa}{}_{\mu\sigma} - \Gamma^{\mu}{}_{\kappa\sigma}\Gamma^{\kappa}{}_{\mu\rho} \\ &= \partial_{\rho}\Gamma^{\mu}{}_{\mu\sigma} - \partial_{\sigma}\Gamma^{\mu}{}_{\mu\rho}. \end{aligned} \quad (98)$$

We have relied so far the $GL(4)$ gauge field only. The other independent field variable, the metric tensor can be used to construct the scalar curvatures, $R = g^{\mu\nu}R_{\mu\nu}$ and $R' = g^{\mu\nu}R'_{\mu\nu}$. The scalar curvature is actually trivial, $R' = 0$, because $R'_{\mu\nu}$ is antisymmetric and its contraction with the symmetric $g^{\mu\nu}$ is vanishing. Note that the curvature tensor is different in the presence of torsion and R' becomes non-trivial.

E. Metric admissibility

A natural relation can be established between the two, otherwise independent quantities, the affine connection and the metric in the following manner. Let us consider two vector fields, $u^{\mu}(x)$ and $v^{\mu}(x)$, which are parallel transported along a path $\gamma(s)$,

$$\dot{\gamma}(s)D_{\mu}u = \dot{\gamma}(s)D_{\mu}v = 0. \quad (99)$$

The connection between the symmetric part of the affine connection, $\{\overset{\rho}{\mu\nu}\} = \frac{1}{2}(\Gamma^{\rho}{}_{\nu\mu} + \Gamma^{\rho}{}_{\mu\nu})$, called Christoffel symbol and the metric structure is achieved by imposing the condition that the scalar product of parallel transported vectors is preserved,

$$\dot{\gamma}(s)D_{\mu}u^{\nu}g_{\nu\rho}v^{\rho} = u^{\nu}v^{\rho}\dot{\gamma}(s)D_{\mu}g_{\nu\rho} = 0 \quad (100)$$

which amounts to the covariant equation,

$$Dg = 0, \quad (101)$$

the metric admissibility condition which fixes the torsion free affine connection. In order to find a more explicit form we write down this equation with all indices shown together with the relations

obtained by performing cyclic permutations on the indices,

$$\begin{aligned}
D_\rho g_{\mu\nu} &= \partial_\rho g_{\mu\nu} - g_{\kappa\nu} \Gamma_{\mu\rho}^\kappa - g_{\mu\kappa} \Gamma_{\nu\rho}^\kappa \\
D_\mu g_{\nu\rho} &= \partial_\mu g_{\nu\rho} - g_{\kappa\rho} \Gamma_{\nu\mu}^\kappa - g_{\nu\kappa} \Gamma_{\rho\mu}^\kappa \\
D_\nu g_{\rho\mu} &= \partial_\nu g_{\rho\mu} - g_{\kappa\mu} \Gamma_{\rho\nu}^\kappa - g_{\rho\kappa} \Gamma_{\mu\nu}^\kappa.
\end{aligned} \tag{102}$$

By taking the sum of the last two equations minus the first one arrives at

$$\Gamma_{\rho\mu\nu} + \Gamma_{\rho\nu\mu} = \partial_\mu g_{\nu\rho} + \partial_\nu g_{\rho\mu} - \partial_\rho g_{\mu\nu} \tag{103}$$

with $\Gamma_{\rho\mu\nu} = g_{\rho\kappa} \Gamma_{\mu\nu}^\kappa$ and the relation

$$\left\{ \begin{matrix} \rho \\ \mu\nu \end{matrix} \right\} = \frac{1}{2} g^{\rho\sigma} (\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\sigma\mu} - \partial_\sigma g_{\mu\nu}). \tag{104}$$

The antisymmetric part of the affine connection, the torsion tensor is left free after imposing the metric admissibility condition.

The curvature tensor, given in terms of the metric, occurs frequently in applications. We have

$$\begin{aligned}
R_{\rho\kappa\nu\mu} &= g_{\rho\lambda} \left[\Gamma_{\kappa\mu,\nu}^\lambda + \Gamma_{\sigma\nu}^\lambda \Gamma_{\kappa\mu}^\sigma - (\mu \iff \nu) \right] \\
&= \frac{1}{2} g_{\rho\lambda} \left\{ \left[g^{\lambda\sigma} (g_{\sigma\kappa,\mu} + g_{\mu\sigma,\kappa} - g_{\kappa\mu,\sigma}) \right]_{,\nu} + \Gamma_{\sigma\nu}^\lambda \Gamma_{\kappa\mu}^\sigma - (\mu \iff \nu) \right\} \\
&= \frac{1}{2} g_{\rho\lambda} g_{,\nu}^{\lambda\sigma} (g_{\sigma\kappa,\mu} + g_{\mu\sigma,\kappa} - g_{\kappa\mu,\sigma}) + \frac{1}{2} (g_{\mu\rho,\kappa\nu} - g_{\kappa\mu,\rho\nu}) + g_{\rho\lambda} \Gamma_{\sigma\nu}^\lambda \Gamma_{\kappa\mu}^\sigma - (\mu \iff \nu)
\end{aligned} \tag{105}$$

where the notation $f_{,\mu} = \partial_\mu f$ is used. The relation

$$g_{\rho\lambda} g_{,\nu}^{\lambda\sigma} = -g^{\lambda\sigma} g_{\rho\lambda,\nu} = -g^{\lambda\sigma} (\Gamma_{\nu\rho\lambda} + \Gamma_{\nu\lambda\rho}) \tag{106}$$

allows us to write

$$\begin{aligned}
R_{\rho\kappa\nu\mu} &= -\frac{1}{2} g^{\lambda\sigma} (\Gamma_{\nu\rho\lambda} + \Gamma_{\nu\lambda\rho}) (g_{\sigma\kappa,\mu} + g_{\mu\sigma,\kappa} - g_{\kappa\mu,\sigma}) + \frac{1}{2} (g_{\mu\rho,\kappa\nu} - g_{\kappa\mu,\rho\nu}) + g_{\rho\lambda} \Gamma_{\sigma\nu}^\lambda \Gamma_{\kappa\mu}^\sigma \\
&\quad - (\mu \iff \nu) \\
&= -\Gamma_{\kappa\mu}^\lambda (\Gamma_{\nu\rho\lambda} + \Gamma_{\nu\lambda\rho}) + \frac{1}{2} (g_{\mu\rho,\kappa\nu} - g_{\kappa\mu,\rho\nu}) + g_{\rho\lambda} \Gamma_{\sigma\nu}^\lambda \Gamma_{\kappa\mu}^\sigma - (\mu \iff \nu) \\
&= \frac{1}{2} (g_{\mu\rho,\kappa\nu} - g_{\nu\rho,\kappa\mu} - g_{\kappa\mu,\rho\nu} + g_{\kappa\nu,\rho\mu}) - g_{\sigma\lambda} \Gamma_{\kappa\mu}^\lambda \Gamma_{\rho\nu}^\sigma - g_{\sigma\rho} \Gamma_{\kappa\mu}^\lambda \Gamma_{\lambda\nu}^\sigma + g_{\rho\lambda} \Gamma_{\sigma\nu}^\lambda \Gamma_{\kappa\mu}^\sigma \\
&\quad + g_{\sigma\lambda} \Gamma_{\kappa\nu}^\lambda \Gamma_{\rho\mu}^\sigma + g_{\sigma\rho} \Gamma_{\kappa\nu}^\lambda \Gamma_{\lambda\mu}^\sigma - g_{\rho\lambda} \Gamma_{\sigma\mu}^\lambda \Gamma_{\kappa\nu}^\sigma \\
&= \frac{1}{2} (g_{\mu\rho,\kappa\nu} - g_{\nu\rho,\kappa\mu} - g_{\kappa\mu,\rho\nu} + g_{\kappa\nu,\rho\mu}) - g_{\sigma\lambda} \Gamma_{\kappa\mu}^\lambda \Gamma_{\rho\nu}^\sigma + g_{\sigma\lambda} \Gamma_{\kappa\nu}^\lambda \Gamma_{\rho\mu}^\sigma.
\end{aligned} \tag{107}$$

Let us consider a two-dimensional sphere as a simple example. The invariant length $ds^2 = r^2(d\theta^2 + \sin^2\theta d\phi^2)$ gives the metric tensor

$$g_{\mu\nu} = r^2 \begin{pmatrix} 1 & 0 \\ 0 & \sin^2\theta \end{pmatrix}, \quad g^{\mu\nu} = \frac{1}{r^2} \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{\sin^2\theta} \end{pmatrix}. \tag{108}$$

The non-vanishing matrix elements of the Christoffel symbol, $\left\{ \begin{smallmatrix} \theta \\ \phi\phi \end{smallmatrix} \right\} = -\sin\theta \cos\theta$, $\left\{ \begin{smallmatrix} \phi \\ \theta\phi \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} \phi \\ \phi\theta \end{smallmatrix} \right\} = -\cot\theta$, give rise

$$\begin{aligned} R_{\phi\theta\phi}^{\theta} &= \partial_{\theta} \left\{ \begin{smallmatrix} \theta \\ \phi\phi \end{smallmatrix} \right\} - \partial_{\phi} \left\{ \begin{smallmatrix} \theta \\ \theta\phi \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} \theta \\ \theta\mu \end{smallmatrix} \right\} \left\{ \begin{smallmatrix} \mu \\ \phi\phi \end{smallmatrix} \right\} - \left\{ \begin{smallmatrix} \theta \\ \phi\mu \end{smallmatrix} \right\} \left\{ \begin{smallmatrix} \mu \\ \theta\phi \end{smallmatrix} \right\} \\ &= -\partial_{\theta} \sin\theta \cos\theta + \sin\theta \cos\theta \cot\theta = \sin^2\theta \end{aligned} \quad (109)$$

and $R_{\theta\theta\phi}^{\phi} = g^{\phi\phi} R_{\phi\theta\theta\phi} = -g^{\phi\phi} R_{\theta\phi\theta\phi} = -g^{\phi\phi} g_{\theta\theta} R_{\phi\theta\phi}^{\theta} = -1$. The Ricci tensor is diagonal, $R_{\theta\theta} = 1$, $R_{\phi\phi} = \sin^2\theta$ and the scalar curvature is $R = \frac{2}{r^2}$.

The symmetries of the curvature tensor in addition to (91) and (93) in the metric admissible case are

$$R_{\rho\kappa\mu\nu} = -R_{\kappa\rho\mu\nu} = R_{\mu\nu\rho\kappa} \quad (110)$$

and the number of independent components is $256 \rightarrow 20$. We note that the curvature is vanishing for flat space only.

The contraction of two indices in the Bianchi-identity (96) gives

$$0 = D_{\mu} R_{\kappa\rho} + D_{\nu} R_{\kappa\rho\mu}^{\nu} - D_{\rho} R_{\kappa\mu}. \quad (111)$$

A further contraction of indices κ and μ by means of the metric tensor gives for the metric admissible curvature tensor

$$0 = 2D_{\mu} R_{\rho}^{\mu} - D_{\rho} R \quad (112)$$

which expresses the covariant conservation law for the Einstein tensor

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R, \quad (113)$$

as

$$D_{\mu} G^{\mu}_{\nu} = 0. \quad (114)$$

Einstein's original argument to establish the equation of motion for the space time geometry was based on a trial and error method of finding a covariant equation whose source term is the energy-momentum tensor,

$$X^{\mu\nu} = T^{\mu\nu}. \quad (115)$$

Due to the conservation law, $D_{\mu} T^{\mu\nu} = 0$, we have $D_{\mu} X^{\mu\nu} = 0$. Since $X^{\mu\nu}$ is expected to be linear in the curvature tensor the choice $X = \kappa G$, κ being a constant, is a natural one.

The metric admissibility simplifies the condition, expressing the invariance of the metric tensor under space-time diffeomorphism. In fact, the vanishing of the Lie derivative of the metric tensor,

$$\begin{aligned} 0 = \nabla_w g_{\mu\nu} &= w^\kappa \partial_\kappa g_{\mu\nu} + \partial_\mu w^\kappa g_{\kappa\nu} + \partial_\nu w^\kappa g_{\mu\kappa} \\ &= D_\mu w_\nu + D_\nu w_\mu + 2S_{\nu\kappa}^\rho g_{\rho\mu} w^\kappa \end{aligned} \quad (116)$$

and the diffeomorphism $w^\mu(x)$, satisfying this condition for a given metric tensor is called Killing field, representing a symmetry of the metric in question.

We close this section by mentioning the question of invariant integral measure and some useful expressions of tensor analysis.

The metric tensor transforms as

$$g_{\mu\nu} = \frac{\partial x'^\rho}{\partial x^\mu} \frac{\partial x'^\sigma}{\partial x^\nu} g'_{\rho\sigma} \quad (117)$$

during the coordinate change $x^\mu \rightarrow x'^\mu$ and its determinant, $g = \det g_{\mu\nu}$, changes as

$$g = g' \left(\det \frac{\partial x'}{\partial x} \right)^2 \quad (118)$$

showing that the integral measure

$$d_{\text{inv}} x = dx \sqrt{-g} \rightarrow dx' \det \frac{\partial x}{\partial x'} \sqrt{-g'} \det \frac{\partial x'}{\partial x} \quad (119)$$

stays invariant. Thus one speaks of scalar, vector and tensor densities constructed by means of the determinant of the metric tensor as $S\sqrt{-g}$, $v^\mu \sqrt{-g}$, $T^{\mu\nu} \sqrt{-g}$, etc.

A useful and simple expression can be obtained for the divergence of a vector field in the absence of torsion by using the equation

$$\delta g = \frac{\partial g}{\partial g_{\sigma\mu}} \delta g_{\sigma\mu} = g g^{\sigma\mu} \delta g_{\sigma\mu} \quad (120)$$

which holds because $g g^{\sigma\mu}$ is actually the minor corresponding to the matrix element $g_{\sigma\mu}$. This gives $\partial_\nu g = g g^{\sigma\mu} \partial_\nu g_{\sigma\mu}$ allows us to write

$$\Gamma_{\nu\mu}^\mu = \frac{1}{2} g^{\sigma\mu} (\partial_\nu g_{\sigma\mu} + \partial_\mu g_{\nu\sigma} - \partial_\sigma g_{\nu\mu}) = \frac{1}{2} g^{\sigma\mu} \partial_\nu g_{\sigma\mu} = \frac{\partial_\nu g}{2g} \quad (121)$$

and

$$\sqrt{-g} \Gamma_{\nu\mu}^\mu = \sqrt{-g} \frac{\partial_\nu g}{2g} = \partial_\nu \sqrt{-g}. \quad (122)$$

The Gauss' theorem can finally be obtained for the divergence

$$D_\mu v^\mu = \partial_\mu v^\mu + \Gamma_{\nu\mu}^\mu v^\nu = \partial_\mu v^\mu + \frac{\partial_\mu \sqrt{-g}}{\sqrt{-g}} v^\mu = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} v^\mu) \quad (123)$$

as

$$\int dx \sqrt{-g} D_\mu v^\mu = \int dx \partial_\mu (\sqrt{-g} v^\mu) = \int ds_\mu \sqrt{-g} v^\mu. \quad (124)$$

A particularly useful application of this relation is for covariantly conserved currents, $D_\mu j^\mu = 0$ which yield ordinarily conserved current density, $\partial_\mu (\sqrt{-g} j^\mu) = 0$ and conserved charge $Q = \int d^3x \sqrt{-g} j^0$.

Metric admissibility renders the definition of the D'Alambertian unique. In fact, we find

$$D_\mu D^\mu = g^{\mu\nu} D_\mu D_\nu = D_\mu g^{\mu\nu} D_\nu = D_\mu D_\nu g^{\mu\nu}, \quad (125)$$

and its action on a scalar is particularly simple,

$$D_\mu D^\mu \phi = D_\mu \partial^\mu \phi = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} \partial^\mu \phi) \quad (126)$$

As an example consider a d -dimensional Euclidean space, parametrized by polar coordinates,

$$x^j = \begin{pmatrix} r \\ r \cos \phi^1 \\ r \sin \phi^1 \cos \phi^2 \\ r \sin \phi^1 \sin \phi^2 \cos \phi^3 \\ \vdots \\ r \sin \phi^1 \dots \sin \phi^{n-2} \cos \phi^{n-1} \\ r \sin \phi^1 \dots \sin \phi^{n-2} \sin \phi^{n-1} \end{pmatrix}, \quad (127)$$

$0 \leq \phi^j \leq \pi$, $j = 1, \dots, d-2$, $0 \leq \phi^{d-1} \leq 2\pi$ where the metric tensor is of the form

$$g_{jk} = r^2 \begin{pmatrix} 1 & 0 \\ 0 & g_{S^{d-1}} \end{pmatrix}. \quad (128)$$

and

$$D^2 = \frac{1}{r^{d-1}} \partial_r r^{d-1} \partial_r + \frac{1}{r^2} D_{S^{d-1}}^2. \quad (129)$$

Another example, the Laplace operator on the two-sphere with metric (108) is

$$\Delta_{S^2} = \frac{1}{r^2} \left[\frac{1}{\sin \theta} \partial_\theta \sin \theta \partial_\theta + \frac{1}{\sin^2 \theta} \partial_\phi^2 \right]. \quad (130)$$

The relation between the affine connection and the metric tensor, the metric admissibility, was imposed as a reasonable assumption. We shall see below that General Relativity supports this relation and the metric admissibility will be the result of the equation of motion for the affine connection.

F. Dynamics

The equations of motion of Einstein's general relativity can be obtained as the Euler-Lagrange equations of the Einstein action

$$S_E = -\frac{1}{16\pi G} \int dx \sqrt{-g} (R + 2\Lambda) = -\frac{1}{16\pi G} \int dx \sqrt{-g} (g^{\mu\nu} R_{\mu\nu} + 2\Lambda) \quad (131)$$

with G and Λ being the gravitational and the cosmological constants. The action is considered as the functional of the independent field variables g and Γ .

The affine connection is not a tensor and it will be advantageous to use tensor fields as independent variables. Thus we separate the Christoffel symbol from the affine connection by writing

$$\Gamma_{\mu\nu}^\rho = \{\}_{\mu\nu}^\rho + C_{\mu\nu}^\rho \quad (132)$$

and consider $C_{\mu\nu}^\rho$, a tensor, rather than the whole $\Gamma_{\mu\nu}^\rho$ as the independent variable controlling the affine structure.

Notice the following slight complication about the choice of the independent components of the metric tensor used for the variational procedure. The derivative of the relation $\delta_\rho^\mu = g^{\mu\nu} g_{\nu\rho}$ with respect to the space-time coordinates gives

$$\begin{aligned} 0 &= \delta g^{\mu\nu} g_{\nu\rho} + g^{\mu\nu} \delta g_{\nu\rho} \\ \delta g_{\mu\nu} &= -g_{\mu\rho} g_{\nu\sigma} \delta g^{\rho\sigma} \end{aligned} \quad (133)$$

thus the metric can not be used in this calculation to change the position of the indices. Instead, we keep $g^{\mu\nu}$ as the independent tensor field for the metric.

A notation which will serve us is the metric admissible covariant derivative,

$$\tilde{D}_\mu v^\nu = \partial_\mu v^\nu + \{\}_{\rho\mu}^\nu v^\rho \quad (134)$$

which is independent of the tensor field $C_{\mu\nu}^\rho$.

We can now write the $GL(4)$ field strength tensor, the Riemann curvature tensor, expressed in term of the 'background field' covariant derivative \tilde{D} as

$$F_{\mu\nu} = [D_\mu, D_\nu] = [\tilde{D}_\mu + C_\mu, \tilde{D}_\nu + C_\nu]. \quad (135)$$

The variation when $C \rightarrow C + \delta C$ generates

$$\begin{aligned} \delta F_{\mu\nu} &= [D_\mu + \delta C_\mu, D_\nu + \delta C_\nu] - [D_\mu, D_\nu] \\ &= (D_\mu + \delta C_\mu)(D_\nu + \delta C_\nu) - (D_\nu + \delta C_\nu)(D_\mu + \delta C_\mu) - D_\mu D_\nu + D_\nu D_\mu \\ &= D_\mu \delta C_\nu + \delta C_\mu D_\nu - D_\nu \delta C_\mu - \delta C_\nu D_\mu + \mathcal{O}(\delta C^2) \\ &= (D_\mu \delta C_\nu) - (D_\nu \delta C_\mu) + \mathcal{O}(\delta C^2) \end{aligned} \quad (136)$$

which gives

$$\delta R^\mu_{\nu\rho\sigma} = D_\rho \delta C^\mu_{\nu\sigma} - D_\sigma \delta C^\mu_{\nu\rho} \quad (137)$$

when the indices are written explicitly and

$$\delta R_{\nu\sigma} = D_\rho \delta C^\rho_{\nu\sigma} - D_\sigma \delta C^\rho_{\nu\rho}. \quad (138)$$

The variation of the curvature scalar, $g^{\nu\sigma} \delta R_{\nu\sigma}$ is a sum of terms which are proportional to δC or $\partial\delta C$. It is a scalar therefore the terms proportional to $\partial\delta C$ can be extended the replacement $\partial\delta C \rightarrow \tilde{D}\delta C$ by the expense of modifying the terms proportional to δC . The result is the expression

$$g^{\nu\sigma} \delta R_{\nu\sigma} = \tilde{D}_\rho v^\rho + \delta C^\kappa_{\nu\rho} K_\kappa^{\nu\rho}, \quad (139)$$

K being linear in C . The actual calculation

$$\begin{aligned} g^{\nu\sigma} \delta R_{\nu\sigma} &= g^{\nu\sigma} (\tilde{D}_\rho \delta C^\rho_{\nu\sigma} - \tilde{D}_\sigma \delta C^\rho_{\nu\rho}) + C_{\kappa\rho}^\rho \delta C^\kappa_{\nu}{}^\nu - \delta C^{\rho}{}^\nu C_{\nu\rho}^\kappa - \delta C^{\rho}{}_{\nu\kappa} C^{\kappa\nu}{}_\rho - C^{\rho}{}_\kappa{}^\nu \delta C^\kappa_{\nu\rho} + \delta C^{\rho}{}_{\kappa\rho} C^{\kappa\nu}{}^\nu + \delta C^{\rho}{}_{\nu\kappa} C^{\kappa\rho}{}^\nu \\ &= \tilde{D}_\rho v^\rho + C_{\kappa\rho}^\rho \delta C^\kappa_{\nu}{}^\nu + \delta C^{\rho}{}_{\kappa\rho} C^{\kappa\nu}{}^\nu - \delta C^{\rho}{}_\kappa{}^\nu C_{\nu\rho}^\kappa - \delta C^{\rho}{}_{\nu\kappa} C^{\kappa\nu}{}_\rho - C^{\rho}{}_\kappa{}^\nu \delta C^\kappa_{\nu\rho} + \delta C^{\rho}{}_{\nu\kappa} C^{\kappa\rho}{}^\nu \\ &= \tilde{D}_\rho v^\rho + C_{\kappa\rho}^\rho \delta C^\kappa_{\nu}{}^\nu + \delta C^{\rho}{}_{\kappa\rho} C^{\kappa\nu}{}^\nu - \delta C^{\kappa}{}_\nu{}^\rho C_{\rho\kappa}^\nu - \delta C^{\kappa}{}_{\nu\rho} C^{\rho\nu}{}_\kappa - C^{\rho}{}_\kappa{}^\nu \delta C^\kappa_{\nu\rho} + \delta C^{\kappa}{}_{\nu\rho} C^{\rho\nu}{}_\kappa \\ &= \tilde{D}_\rho v^\rho - \tilde{D}_\sigma (g^{\nu\sigma} \delta C^\rho_{\nu\rho}) + \delta C^{\kappa}{}_\nu{}^\rho C_{\kappa\rho}^\rho + \delta C^{\rho}{}_{\kappa\rho} C^{\kappa\nu}{}^\nu - \delta C^{\kappa}{}_{\nu\rho} (C^{\nu\rho}{}_\kappa - C^{\rho\nu}{}_\kappa - C^{\rho}{}_\kappa{}^\nu + C^{\rho\nu}{}_\kappa) \\ &= \tilde{D}_\rho v^\rho + \delta C^{\kappa}{}_{\nu\rho} (g^{\nu\rho} C_{\kappa\rho}^\rho + g^{\kappa\rho} C_{\lambda}^{\nu\lambda} - C^{\nu\rho}{}_\kappa + C^{\rho\nu}{}_\kappa) \end{aligned} \quad (140)$$

yields

$$v^\rho = g^{\nu\sigma} \delta C^\rho_{\nu\sigma} - g^{\nu\rho} \delta C^\sigma_{\nu\sigma}, \quad (141)$$

and

$$K_\kappa^{\nu\rho} = g^{\nu\rho} C_{\kappa\rho}^\rho + g^{\kappa\rho} C_{\lambda}^{\nu\lambda} - C^{\nu\rho}{}_\kappa + C^{\rho\nu}{}_\kappa. \quad (142)$$

After these preparation one can easily calculate the variation of the integrand in the action

$$\delta[\sqrt{-g}(R_{\mu\nu}g^{\mu\nu} + 2\Lambda)] = \delta\sqrt{-g}(R_{\mu\nu}g^{\mu\nu} + 2\Lambda) + \sqrt{-g}\delta R_{\mu\nu}g^{\mu\nu} + \sqrt{-g}R_{\mu\nu}\delta g^{\mu\nu}. \quad (143)$$

The first term contains

$$\begin{aligned} \delta\sqrt{-g} &= -\frac{g}{2\sqrt{-g}}g^{\mu\nu}\delta g_{\mu\nu} \\ &= -\frac{1}{2}\sqrt{-g}g^{\mu\nu}g_{\mu\rho}g_{\nu\sigma}\delta g^{\rho\sigma} \\ &= -\frac{1}{2}\sqrt{-g}g_{\nu\sigma}\delta g^{\nu\sigma}, \end{aligned} \quad (144)$$

the second is given by Eq. (139) thus we have

$$\begin{aligned} \delta[\sqrt{-g}(R_{\mu\nu}g^{\mu\nu} + 2\Lambda)] &= -\frac{1}{2}\sqrt{-g}g_{\nu\sigma}\delta g^{\nu\sigma}(R + 2\Lambda) + \sqrt{-g}\tilde{D}_\rho v^\rho + \sqrt{-g}\delta C^\kappa_{\nu\rho}K_\kappa^{\nu\rho} + \sqrt{-g}R_{\mu\nu}\delta g^{\mu\nu} \\ &= \sqrt{-g}\left(R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R - \Lambda g_{\mu\nu}\right)\delta g^{\mu\nu} + \sqrt{-g}\delta C^\kappa_{\nu\rho}K_\kappa^{\nu\rho} + \sqrt{-g}\tilde{D}_\rho v^\rho \end{aligned} \quad (145)$$

The last contribution can be ignored being a surface term which is ignored from the point of view of the equations of motion. The variation of the affine connection yields $K = 0$ which gives $C = 0$, the metric admissibility condition for the affine connection. Finally, the variation of the metric tensor leads to the Einstein equation

$$G_{\mu\nu} - \Lambda g_{\mu\nu} = 0 \quad (146)$$

in the absence of matter.

IV. COUPLING TO MATTER

After having obtained the Lagrangian and the equations of motion for the gravitation field in the absence of matter let us turn to the question of introducing matter in our description.

A. Point particle in an external gravitational field

First we consider the simplest problem of the motion of a massive point particle in a fixed gravitational field, ie. in a fixed geometry where the trajectory, identified by the equation of motion is the generalization of the straight line of the Minkowskian, flat space-time.

1. Equivalence Principle

The simplest way to find the equation of motion for a point particle is to use the Equivalence principle. A free particle follows the straight trajectory $\xi^a(s)$ satisfying the equation of motion

$$\frac{d\dot{\xi}^a(s)}{ds} = 0 \quad (147)$$

with $\dot{\xi} = \frac{d\xi}{ds}$ in flat space-time, in the absence of gravity. When an external gravitational field is introduced then the trajectory $x^\mu(s)$ is not a straight line anymore but the Equivalence Principle allows us to recover the same equation of motion locally, at a given space-time point, by an appropriate choice of the coordinate system. Eq. (147) shows that the four velocity, assumed as a vector field, $u^\mu(x) = \dot{x}^\mu$, filling up a region of the space-time, remains unchanged on the world line. The unique covariant extension of such a parallel transport on a non-flat geometry is

$$u^\nu D_\nu u^\mu = \dot{u}^\mu + u^\rho \Gamma^\mu_{\rho\nu} u^\nu = 0. \quad (148)$$

Let us now assume that an external, non-gravitational force acts on the particle and Eq. (148) is replaced by

$$\dot{u}^\mu + u^\rho \Gamma_{\rho\nu}^\mu u^\nu = \frac{F^\mu}{mc}. \quad (149)$$

The form

$$mc\dot{u}^\mu = F^\mu + F_{gr}^\mu \quad (150)$$

of this equation with

$$F_{gr}^\mu = -mu^\mu \Gamma_{\rho\nu}^\mu u^\nu \quad (151)$$

shows that the gravitational field generates a force F_{gr} which is linear in the velocities in a manner similar to the Lorentz force of electrodynamics where

$$\begin{aligned} mc\ddot{x}^\mu &= F^\mu + F_{ed}^\mu \\ F_{ed}^\mu &= \frac{e}{c} F_{\nu}^\mu u^\nu, \end{aligned} \quad (152)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the field strength tensor.

2. Spin precession

It is instructive to generalize this argument for a gyroscope, a point particle with an angular momentum, spin, given by the four-vector $S^\mu = (0, \mathbf{S})$ in the rest-frame. In the absence of external forces we have time independent spin,

$$\frac{dS^\mu}{ds} = 0. \quad (153)$$

The covariant generalization for an external gravitational field of the form $S^\mu = (0, \mathbf{S})$ of the spin vector is the equation of motion

$$\dot{S}^\mu + \Gamma_{\rho\nu}^\mu S^\nu \dot{x}^\rho = 0, \quad (154)$$

together with the auxiliary condition

$$S_\mu \dot{x}^\mu = 0. \quad (155)$$

We furthermore assume that the external force F_{ext} does not exert torque on the system. The spin will still be conserved, $d\mathbf{S}/dt = 0$, in the co-moving frame which can be reproduced in a covariant form by requiring that the vector \dot{S} be proportional to \dot{x} ,

$$\dot{S} = a\dot{x}. \quad (156)$$

The covariant derivative of the orthogonality condition (155) along the world line, $0 = a\dot{x}_\mu\dot{x}^\mu + S_\mu\ddot{x}^\mu$, yields

$$a = -S_\mu\ddot{x}^\mu = -S_\mu\frac{F^\mu}{m}. \quad (157)$$

The covariant generalization of the equation of motion (156),

$$\dot{S}^\mu = -S_\nu\frac{F^\nu}{m}\dot{x}^\mu, \quad (158)$$

is the Fermi-Walker transport of the spin which reduces to parallel transport in the absence of external force, $F = 0$.

3. Variational equation of motion

A direct derivation of the trajectory of a point particle, without referring to the Equivalence Principle starts with the action of a free massive point particle,

$$S = -mc \int \sqrt{\dot{x}^\mu g_{\mu\nu}(x)\dot{x}^\nu} d\tau \quad (159)$$

where $\dot{x}^\mu = dx^\mu(\tau)/d\tau$. The corresponding Euler-Lagrange equation,

$$\frac{\partial L}{\partial x^\rho} - \frac{d}{d\tau} \frac{\partial L}{\partial \dot{x}^\rho} = 0 \quad (160)$$

containing the terms

$$\begin{aligned} \frac{\partial L}{\partial x^\rho} &= -mc \frac{\dot{x}^\mu \partial_\rho g_{\mu\nu} \dot{x}^\nu}{2\sqrt{\dot{x}^\mu g_{\mu\nu} \dot{x}^\nu}}, \\ \frac{\partial L}{\partial \frac{dx^\rho}{d\tau}} &= -mc \frac{g_{\rho\nu} \dot{x}^\nu}{\sqrt{\dot{x}^\mu g_{\mu\nu} \dot{x}^\nu}}, \end{aligned} \quad (161)$$

reads

$$\begin{aligned} 0 &= -\frac{\dot{x}^\mu \partial_\rho g_{\mu\nu} \dot{x}^\nu}{2\sqrt{\dot{x}^\mu g_{\mu\nu} \dot{x}^\nu}} + \frac{d}{d\tau} \frac{g_{\rho\nu} \dot{x}^\nu}{\sqrt{\dot{x}^\mu g_{\mu\nu} \dot{x}^\nu}} \\ &= \frac{1}{\sqrt{\dot{x}^\mu g_{\mu\nu} \dot{x}^\nu}} \left[-\frac{1}{2} \dot{x}^\mu \partial_\rho g_{\mu\nu} \dot{x}^\nu + \dot{x}^\kappa \partial_\kappa g_{\rho\nu} \dot{x}^\nu + g_{\rho\nu} \ddot{x}^\nu + g_{\rho\nu} \dot{x}^\nu \frac{d}{d\tau} \frac{1}{\sqrt{\dot{x}^\mu g_{\mu\lambda} \dot{x}^\lambda}} \right] \end{aligned} \quad (162)$$

We symmetrize in the indices κ and ν the factor $\partial_\kappa g_{\rho\nu}$ in the second term of the last line and find

$$\begin{aligned} 0 &= -\frac{1}{2} \dot{x}^\mu \partial_\rho g_{\mu\nu} \dot{x}^\nu + \frac{1}{2} \dot{x}^\kappa (\partial_\kappa g_{\rho\nu} + \partial_\nu g_{\rho\kappa}) \dot{x}^\nu + g_{\rho\nu} \ddot{x}^\nu + g_{\rho\nu} \frac{dx^\nu}{d\tau} \frac{d}{d\tau} \frac{1}{\sqrt{\dot{x}^\mu g_{\mu\lambda} \dot{x}^\lambda}} \\ &= g_{\rho\sigma} (\ddot{x}^\sigma + \Gamma_{\nu\kappa}^\sigma \dot{x}^\nu \dot{x}^\kappa + \dot{x}^\sigma \dot{f}) \end{aligned} \quad (163)$$

with $f(\tau) = 1/\sqrt{\dot{x}^\mu g_{\mu\lambda} \dot{x}^\lambda}$. This equation assumes the simplest form (148) when $f(\tau)$ is constant, ie. $\tau = s$, the invariant length of the world line and its solution is called geodesic. For $\tau \neq s$ the term, proportional to \dot{f} adjusts the length of the four-velocity without changing its direction.

4. Geodesic deviation

The dynamics of an infinitesimal deviation from a solution of the Newton equation of a particle moving in a given potential is that of a harmonic oscillator with time dependent frequency. In fact, let us consider a trajectory $\xi(t)$ which obeys the equation of motion, $m\ddot{\xi}^j = -\nabla^j \mathcal{V}$ and consider a neighboring trajectory $\xi(t) + \delta\xi(t)$. The infinitesimal shift $\delta\mathbf{x}$ satisfies the linear, time dependent equation $m\delta\ddot{\xi}^j = -\delta\xi^k \nabla^k \nabla^j \mathcal{V}$.

The generalization of the equation of motion of infinitesimal deviation $\delta\xi^\mu$ from a geodesic consists of a straightforward linearization of (148) in the deformation. But the result is easier to find by embedding our world line and its deformed partner into a family of world lines, solutions of Eq. (148) which fill up the space-time in the vicinity of our observation point and using a curve $\gamma^\mu(\tau)$ which crosses our world line and its deformed partner. The integral curves of the four-velocity vector field, $u(x) = \frac{d}{ds}\xi(x) = \dot{\xi}(x)$, are the world lines, hence $\frac{d}{ds}\phi = u^\nu D_\nu \phi = \dot{\phi}$ holds for any field $\phi(x)$.

Let us consider the surface in the space-time which is swept through by the world lines which cross γ and use the coordinates s and τ to identify its points $\xi^\mu(s, \tau)$. The coordinate basis vector fields, $u = \partial_s \xi(s, \tau)$, $v = \partial_\tau \xi(s, \tau)$ are holonomic, $\partial_s v = \partial_\tau u$ according to Eq. (63), and we have

$$u^\nu D_\nu v^\mu = v^\nu D_\nu u^\mu. \quad (164)$$

Let us calculate finally the acceleration of the deformation $\delta\xi = \epsilon \partial_\tau \xi = \epsilon v$,

$$\ddot{v} = u^\mu D_\mu (u^\nu D_\nu v) \quad (165)$$

Two successive applications of the holonomy condition give

$$\begin{aligned} \ddot{v} &= u^\mu D_\mu (v^\nu D_\nu u) \\ &= u^\mu (D_\mu v^\nu) D_\nu u + u^\mu v^\nu D_\mu D_\nu u \\ &= v^\mu (D_\mu u^\nu) D_\nu u + u^\mu v^\nu [D_\mu, D_\nu] u + u^\mu v^\nu D_\nu D_\mu u \\ &= v^\mu (D_\mu u^\nu) D_\nu u + u^\mu v^\nu [D_\mu, D_\nu] u + v^\nu D_\nu (u^\mu D_\mu u) - v^\nu (D_\nu u^\mu) D_\mu u \\ &= u^\mu v^\nu [D_\mu, D_\nu] u + v^\nu D_\nu (u^\mu D_\mu u) \end{aligned} \quad (166)$$

The four-acceleration of the world lines is vanishing, therefore

$$\ddot{v}^\rho = R^\rho_{\ \kappa\mu\nu} u^\kappa u^\mu v^\nu, \quad (167)$$

the acceleration satisfies a linear equation whose coefficient matrix depends is a quadratic function of the four-velocity.

It is instructive to see what happens in electrodynamics where we may start with the the equation of motion, (152), written for the deformed world line $x + \delta x$ is

$$mc(\ddot{x}^\mu + \delta\ddot{x}^\mu) = \frac{e}{c}F^\mu{}_\nu(x + \delta x)(\dot{x}^\nu + \delta\dot{x}^\nu). \quad (168)$$

The linearization in the deformation δx yields immediately

$$mc\delta\ddot{x}^\mu = \frac{e}{c}\delta x^\rho\partial_\rho F^\mu{}_\nu\dot{x}^\nu + \frac{e}{c}F^\mu{}_\nu\delta\dot{x}^\nu. \quad (169)$$

This equation can be obtained from Eq. (166) by first making the replacement $D_\mu \rightarrow \partial_\mu$,

$$mc\ddot{v} = v^\nu\partial_\nu\dot{u}, \quad (170)$$

followed by the use of the equation of motion, Eq. (152),

$$mc\ddot{v}^\mu = \frac{e}{c}v^\nu\partial_\nu F^\mu{}_\rho u^\rho + \frac{e}{c}F^\mu{}_\rho v^\nu\partial_\nu u^\rho. \quad (171)$$

Eq. (169) follows by noting $v^\nu\partial_\nu u^\rho = \delta x^\nu\partial_\nu\dot{x}^\rho = \delta\tau\partial_\tau\partial_s x^\rho = \partial_s(\delta\tau\partial_\tau x^\rho) = \delta\dot{x}^\rho$.

5. Newtonian limit

It is instructive to consider the Newtonian limit where the static gravitational field is weak and the motion of the test particle is slow by writing

$$g_{\mu\nu} = \eta_{\mu\nu} + \gamma_{\mu\nu} \quad (172)$$

and assuming that γ is an infinitesimal tensor. The slow motion of the test particle leads to $\frac{dx^\mu}{ds} \approx (1, 0, 0, 0)$ which together with $g^{\mu\nu} = \eta^{\mu\nu} - \gamma^{\mu\nu}$ gives

$$\ddot{x}^\mu \approx -\Gamma_{00}^\mu = \frac{1}{2}\frac{\partial\gamma_{00}}{\partial x^\mu}, \quad (173)$$

or

$$\ddot{\mathbf{x}} = -\nabla\phi \quad (174)$$

where the static Newtonian potential is

$$\phi = -\frac{\gamma_{00}}{2}. \quad (175)$$

B. Interacting matter-gravity system

The variation of the the Einstein action S_E with respect to the metric tensor leads to the vacuum Einstein equation, Eq. (146). When matter is included then the action becomes the sum of the gravitational and the matter actions,

$$S = S_E + S_M. \quad (176)$$

Therefore the matter contribution to the Einstein equation will be given by the expression

$$G_{\mu\nu} - \Lambda g_{\mu\nu} = 8\pi G M_{\mu\nu}(x) \quad (177)$$

with

$$M_{\mu\nu}(x) = \frac{2}{\sqrt{-g}} \frac{\delta S_M}{\delta g^{\mu\nu}(x)} \quad (178)$$

The source for gravitational interaction is the mass. According to special relativity this corresponds to energy and covariance makes the whole energy-momentum tensor as the source. Thus one expects that the quantity (178) is proportional to the energy momentum tensor. But the Einstein equation (177) then expresses the vanishing of the full energy-momentum tensor. The vanishing of the total energy-momentum tensor is understandable because this latter is defined by carrying out space-time translations, an operation which becomes ill-defined without a background space-time.

A nontrivial condition, satisfied by the Einstein equation in the presence of matter is Eq. (114), amounts to the energy-momentum conservation, $D_\mu M^\mu{}_\nu = 0$, for any theory with action of the form (176). Hence the energy-momentum of the matter and the gravitation field are conserved separately. It is remarkable is that the gravity-matter interaction does not violate the conservation of the matter energy-momentum and the energy-momentum density and flux, the matrix elements of the energy-momentum tensor, are identical for the gravity and mater, except their sign.

We review briefly the energy-momentum tensor of a system of point particles, ideal fluid and a scalar field.

1. Point particle

Let us suppose that we have a particle of mass m moving along the world lines x^μ . The action is

$$S_M = -mc \int ds \sqrt{\dot{x}^\mu g_{\mu\nu} \dot{x}^\nu} \quad (179)$$

whose variation

$$\begin{aligned}
\delta S_M &= -\frac{1}{2}mc \int ds \frac{\dot{x}^\mu \delta g_{\mu\nu} \dot{x}^\nu}{\sqrt{\dot{x}^\mu g_{\mu\nu} \dot{x}^\nu}} \\
&= -\frac{1}{2}mc \int ds \dot{x}^\mu \dot{x}^\nu \delta g_{\mu\nu} \\
&= -\frac{1}{2} \int dx \sqrt{-g} \int ds \frac{p^\mu(s(t)) p^\nu(s(t))}{mc} \delta(x - x(s)) \delta g_{\mu\nu}(x),
\end{aligned} \tag{180}$$

where the parameter of the world line is chosen to be the invariant length in the second equation, after having completed the variation. The definition (178) yields

$$M^{\mu\nu}(x) = \int ds \frac{p^\mu(s(t)) p^\nu(s(t))}{mc} \delta(x - x(s)). \tag{181}$$

The density of the four-momentum is

$$T^{\mu 0}(t, \mathbf{x}) = p^\mu(s(t)) \delta(\mathbf{x} - \mathbf{x}(t)). \tag{182}$$

The tensor which reduces to this expression is

$$T^{\mu\nu}(x) = \frac{p^\mu(s(t)) p^\nu(s(t))}{p^0(s(t))} \delta(\mathbf{x} - \mathbf{x}(t)), \tag{183}$$

and it can be written in a manifestly covariant form as

$$\begin{aligned}
T^{\mu\nu}(x) &= \int ds \frac{p^\mu(s(t)) p^\nu(s(t))}{p^0(s(t))} c \frac{dt}{ds} \delta(x - x(s)) \\
&= \int ds \frac{p^\mu(s(t)) p^\nu(s(t))}{mc} \delta(x - x(s)),
\end{aligned} \tag{184}$$

establishing $M = T$.

2. Ideal fluid

It is worthwhile mentioning that in a more realistic situation one assumes a continuous distribution of matter. For homogeneous and isotropic matter in the rest frame we have

$$T^{\mu\nu} = \begin{pmatrix} \epsilon & 0 & 0 & 0 \\ 0 & p & 0 & 0 \\ 0 & 0 & p & 0 \\ 0 & 0 & 0 & p \end{pmatrix} \tag{185}$$

which can be written in a covariant manner as

$$T^{\mu\nu} = (p + \epsilon) \dot{x}^\mu \dot{x}^\nu - p g^{\mu\nu}, \tag{186}$$

because the relation $\dot{x}^\mu = (1, 0, 0, 0)$ holds in the rest frame. For ideal fluid where mean free path and times are short enough to maintain isotropy we have

$$T^{\mu\nu}(x) = (p(x) + \epsilon(x))u^\mu(x)u^\nu(x) - p(x)g^{\mu\nu}, \quad (187)$$

as the source term to the Einstein equation where $u^\mu(x) = \dot{x}^\mu(x)$ is the four-velocity of the fluid particles at the space-time point x .

3. Classical fields

We consider finally a simple scalar field theory with the action

$$S_M = \int dx \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi(x) \partial_\nu \phi(x) - V(\phi(x)) \right]. \quad (188)$$

The calculation of the variation with respect to the metric tensor is greatly simplified by the fact that the Lagrangian depends on the metric tensor and not its space-time derivatives,

$$M_{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\partial \sqrt{-g}}{\partial g^{\mu\nu}} L + 2 \frac{\partial L}{\partial g^{\mu\nu}}. \quad (189)$$

The use of Eq. (120) which gives $g^{\mu\nu}$ as the coefficient of the Lagrangian in the first term and the relation

$$\frac{\partial L}{\partial g^{\mu\nu}} = \frac{1}{2} g_{\nu\kappa} \frac{\partial L}{\partial \partial_\kappa \phi} \partial_\mu \phi \quad (190)$$

establishes the identity of $M_{\mu\nu}$ with the energy-momentum tensor

$$T_{\mu\nu} = \frac{\partial L}{\partial \partial^\mu \phi} \partial_\nu \phi - g_{\mu\nu} L, \quad (191)$$

given by (A44).

V. GRAVITATIONAL RADIATION

A gauge theory has two distinct sectors. One the one hand, there are dynamical degrees of freedom distributed in space-time in such a manner that they support a retarded or advanced signal, generated by external charges, like the far or radiation field in electrodynamics. One the other hand, there are “slave” modes which follow the motion of the external charges in an algebraic manner, without any non-trivial dynamics, like the near or Coulomb field of electrodynamics. We turn now to the former and mention few rudimentary features of gravitational radiation.

Let us start with the state without radiation: the flat Minkowski space-time solves trivially the Einstein equation. A weak radiation field,

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad (192)$$

with $|h_{\mu\nu}| \ll 1$ should not change the space-time geometry in a fundamental manner, an assumption which allows to consider General Relativity on the flat background space-time as a relativistic classical field theory. We shall discuss the plane wave solutions of the linearized Einstein equation in what follows.

A. Linearization

We have up to $\mathcal{O}(h)$

$$\begin{aligned} \Gamma^\rho{}_{\mu\nu} &= \frac{1}{2}\eta^{\rho\sigma}(\partial_\mu h_{\nu\sigma} + \partial_\nu h_{\sigma\mu} - \partial_\sigma h_{\mu\nu}) = \frac{1}{2}(\partial_\mu h_\nu^\rho + \partial_\nu h_\mu^\rho - \partial^\rho h_{\mu\nu}), \\ R^\mu{}_{\nu\rho\sigma} &= \partial_\rho \Gamma^\mu{}_{\nu\sigma} - \partial_\sigma \Gamma^\mu{}_{\nu\rho} = \frac{1}{2}\partial_\rho(\partial_\nu h_\sigma^\mu + \partial_\sigma h_\nu^\mu - \partial^\mu h_{\nu\sigma}) - (\rho \leftrightarrow \sigma), \\ R_{\nu\sigma} &= \frac{1}{2}\partial_\mu(\partial_\nu h_\sigma^\mu + \partial_\sigma h_\nu^\mu - \partial^\mu h_{\nu\sigma}) - \frac{1}{2}\partial_\sigma(\partial_\nu h_\mu^\mu + \partial_\mu h_\nu^\mu - \partial^\mu h_{\nu\mu}) \\ &= \frac{1}{2}(\partial_\nu \partial_\mu h_\sigma^\mu + \partial_\sigma \partial_\mu h_\nu^\mu - \square h_{\nu\sigma} - \partial_\sigma \partial_\nu h), \\ R &= \partial_\nu \partial_\mu h^{\mu\nu} - \square h, \end{aligned} \quad (193)$$

where the indices are raised and lowered by $\eta^{\mu\nu}$ and $\eta_{\mu\nu}$, respectively as in an ordinary relativistic field theory and $h = h^\mu{}_\mu$.

A $GL(4)$ gauge transformation, an external diffeomorphism $x^\mu \rightarrow x^\mu + \xi^\mu(x)$ induces

$$h_{\mu\nu} \rightarrow h'_{\mu\nu} = h_{\mu\nu} + \partial_\mu \xi_\nu + \partial_\nu \xi_\mu. \quad (194)$$

The linearization in $h_{\mu\nu}$ suppresses the commutator of the field strength tensor in the third equation which now looks as an Abelian field strength. As a result our expression for the Riemann, Ricci and the Einstein tensors, as well as the scalar curvature are gauge invariant.

We shall use harmonic gauge (77) where

$$\partial^\nu h_{\mu\nu} = \frac{1}{2}\partial_\mu h. \quad (195)$$

This condition is satisfied after the gauge transformation which solves

$$\square \xi_\nu = \frac{1}{2}\partial_\nu h - \partial_\rho h_\nu^\rho \quad (196)$$

since

$$\frac{\partial}{\partial x^{\nu}} h_{\mu}^{\nu} = \partial^{\nu} (h_{\mu\nu} + \partial_{\mu} \xi_{\nu} + \partial_{\nu} \xi_{\mu}) = \partial^{\nu} h_{\mu\nu} + \partial^{\nu} \partial_{\mu} \xi_{\nu} + \frac{1}{2} \partial_{\mu} h - \partial_{\rho} h_{\mu}^{\rho} = \frac{1}{2} \partial_{\mu} h + \partial_{\mu} \partial^{\nu} \xi_{\nu} = \frac{1}{2} \frac{\partial}{\partial x^{\mu}} h'. \quad (197)$$

Note that the harmonic gauge condition is preserved by further gauge transformations which correspond to harmonic functions, $\square \xi^{\mu} = 0$.

B. Wave equation

The linearized Einstein equation, (177),

$$\frac{1}{2} (\partial_{\nu} \partial_{\mu} h_{\sigma}^{\mu} + \partial_{\sigma} \partial_{\mu} h_{\nu}^{\mu} - \square h_{\nu\sigma} - \partial_{\sigma} \partial_{\nu} h - \eta_{\nu\sigma} \partial_{\rho} \partial_{\mu} h^{\mu\rho} + \eta_{\nu\sigma} \square h) - \Lambda h_{\nu\sigma} = 8\pi G T_{\nu\sigma} \quad (198)$$

contains the $\mathcal{O}(h^0)$, Minkowski energy-momentum tensor because $h = \mathcal{O}(G)$. The wave equation reads in harmonic gauge as

$$\square h_{\nu\sigma} - \frac{1}{2} \eta_{\nu\sigma} \square h + 2\Lambda h_{\nu\sigma} = -16\pi G T_{\nu\sigma} \quad (199)$$

in harmonic gauge.

The equations simplify when expressed in terms of

$$\bar{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} h \quad (200)$$

where $\bar{h}_{\mu}^{\mu} = \bar{h} = -h$ because the gauge condition and the Einstein equation are

$$\partial_{\nu} \bar{h}_{\mu}^{\nu} = 0, \quad (201)$$

and

$$\square \bar{h}_{\nu\sigma} + \Lambda (2\bar{h}_{\nu\sigma} - \eta_{\nu\sigma} \bar{h}) = -16\pi G T_{\nu\sigma}, \quad (202)$$

respectively. The cosmological constant, Λ , plays the role of a mass which makes the radiation field short ranged and will be ignored below. The retarded solution is

$$\bar{h}_{\nu\sigma}(t, \mathbf{x}) = 4G \int d^3 x' \frac{T_{\nu\sigma}(t - |\mathbf{x} - \mathbf{x}'|)}{|\mathbf{x} - \mathbf{x}'|} \quad (203)$$

which is automatically given in harmonic gauge due to $\partial_{\nu} T^{\nu\mu} = 0$.

C. Plane-waves

The linearized Einstein equation in vacuum is satisfied by plane waves,

$$\bar{h}_{\mu\nu} = e_{\mu\nu}e^{ikx} + e_{\mu\nu}^*e^{-ikx}, \quad (204)$$

given in terms of the polarization tensor $e_{\mu\nu}$. The wave vector is light-like, $k^2 = 0$ and the gravitational wave propagates on the line cone. The symmetric polarization tensor contains 10 parameters but the gauge fixing,

$$k^\mu e_{\mu\nu} = 0 \quad (205)$$

decreases the number of the independent parameters to 6.

We may perform a further gauge transformation, $x^\mu \rightarrow x^\mu + \xi^\mu$ with $\square\xi^\mu = 0$ without leaving the harmonic gauge. The choice

$$\xi^\mu = ia^\mu e^{ikx} - ia^{*\mu} e^{-ikx} \quad (206)$$

transforms the polarization tensor into

$$e_{\mu\nu} \rightarrow e'_{\mu\nu} = e_{\mu\nu} + k_\mu a_\nu + k_\nu a_\mu \quad (207)$$

and leaves 2 independent parameters. Some simplification of the covariant expressions can be achieved by choosing the longitudinal and transverse component b^μ in such a manner that $e'_\mu \rightarrow e''_\mu + 2k^\mu a_\mu = 0$ and $e_{\mu 0} \rightarrow e_{\mu 0} + k_\mu a_0 + k_0 a_\mu = 0$, respectively.

It is instructive to compare this situation with electrodynamics in Lorentz gauge, $\partial^\mu A_\mu = 0$, where the plane waves satisfy the wave equation $\square A_\mu = 0$. The solution

$$A_\mu(x) = e_\mu e^{ikx} + e_\mu^* e^{-ikx}, \quad (208)$$

$k^2 = 0$, has 4 independent parameters in the polarization vector e_μ and this number is reduced to 3 by the gauge condition, $k^\mu e_\mu = 0$. The gauge transformation, $A_\mu \rightarrow A_\mu + \partial_\mu \phi$, performed by a harmonic function $\square\phi = 0$ leaves the Lorentz gauge condition unchanged and the choice $\phi(x) = ia e^{ikx} - ia^* e^{-ikx}$ transforms the polarization vector, $e_\mu \rightarrow e_\mu - ak_\mu$, and can be used to reduce the free parameters to 2.

D. Polarization

Let us now consider a simple application, the effect of a gravitational plane wave on the motion of point particles, namely the equation of motion for the 5(t)3.003f

in Section IV A 4. We assume stationary unperturbed particles, $u^\mu = (1, \mathbf{0}) + \mathcal{O}(h)$, $v = \mathcal{O}(h)$ and write Eq. (167) as

$$\partial_0^2 v^\rho = R^\rho_{00\nu} v^\nu = \frac{1}{2} \partial_0^2 h_\nu^\rho v^\nu, \quad (209)$$

where we used $h_{\mu 0} = 0$ in the second equation. The gauge condition (205) shows that the deformation is transverse, $k_\mu v^\mu = 0$. Since $u_\mu v^\mu = 0$ the deformation is transverse in the spatial directions, as well, $\mathbf{k}\mathbf{v} = 0$, with $k^\mu = (k^0, \mathbf{k})$ and $v^\mu = (0, \mathbf{v})$.

We shall use coordinates where $k^\mu = (k, 0, 0, k)$. We know that (i) $k^\mu e_{\mu\nu} = 0$, (ii) $e_\mu^\mu = 0$ and (iii) $e_{0\nu} = 0$. The equations (i) and (iii) imply $e_{3\nu} = 0$, leaving $e_{jk} \neq 0$ for $j, k = 1, 2$ only. Due to (ii) the symmetric matrix $e_{\mu\nu}$ is traceless, leaving two independent components, $e_{11} = -e_{22}$ and $e_{12} = e_{21}$. When $e_{12} = 0$ one finds

$$\partial_0^2 v^j = -\frac{1}{2} k^{02} (e_{11} e^{ikx} + e_{11}^* e^{-ikx}) \epsilon^{jk} v^k, \quad (210)$$

yielding the solution

$$\begin{pmatrix} v^1(x) \\ v^2(x) \end{pmatrix} = \begin{pmatrix} [1 + \frac{1}{2}(e_{11} e^{ikx} + e_{11}^* e^{-ikx})] v^1(0, \mathbf{x}) \\ [1 - \frac{1}{2}(e_{11} e^{ikx} + e_{11}^* e^{-ikx})] v^2(0, \mathbf{x}) \end{pmatrix}. \quad (211)$$

A ring of particle in the (1, 2) plane oscillates horizontally and vertically, as shown in Fig. 1 (a). For $e_{11} = 0$ we have

$$\partial_0^2 v^2 = -\frac{1}{2} k^{02} (e_{12} e^{ikx} + e_{12}^* e^{-ikx}) v^1, \quad (212)$$

and

$$\begin{pmatrix} v^1(x) \\ v^2(x) \end{pmatrix} = \begin{pmatrix} [1 + \frac{1}{2}(e_{12} e^{ikx} + e_{12}^* e^{-ikx})] v^2(0, \mathbf{x}) \\ [1 + \frac{1}{2}(e_{12} e^{ikx} + e_{12}^* e^{-ikx})] v^1(0, \mathbf{x}) \end{pmatrix}. \quad (213)$$

The direction of the deformation of the ring in the (1, 2) plane rotated by $\pi/4$ compared to the previous case as shown in Fig. 1 (b). The direction independent, isotropic deformation is due to the monopole term of the multipole expansion, the deformation along a fixed direction indicate the presence of the dipole term and these deformations, carried out in two directions belong to the quadrupole order.

In the case of electrodynamics the plane wave (208) corresponds to the field strength tensor

$$F_{\mu\nu} = ik_\mu (e_\nu e^{ikx} - e_\nu^* e^{-ikx}) - (\mu \leftrightarrow \nu), \quad (214)$$




FIG. 1: The shape of a circle of particles in the (1,2) plane as a function of time for (a): $e_{12} = 0$ (b): $e_{12} = 0$.

which gives for Eq. (169)

$$\begin{aligned}
 mc\delta\ddot{x}^\mu = & \frac{e}{c}\{i[k^\mu(e_\nu e^{ikx} - e_\nu^* e^{-ikx}) - k_\nu(e^\mu e^{ikx} - e^{*\mu} e^{-ikx})]\delta\dot{x}^\nu \\
 & - \delta x^\rho k_\rho [k^\mu(e_\nu e^{ikx} + e_\nu^* e^{-ikx}) - k_\nu(e^\mu e^{ikx} + e^{*\mu} e^{-ikx})]\dot{x}^\nu\}. \quad (215)
 \end{aligned}$$

A plane wave, propagating in the spatial 3 direction is polarized in the (1,2) plane and one finds deformations, characteristic of dipole field.

Gravitational radiation is too weak to be seen in a direct manner. But an indirect evidence is known, the slowing of the PSR1913+16 binary system, pulsar, is consistent with the energy loss, caused by the power, radiated.

VI. SCHWARZSCHILD SOLUTION

After a short digression into the dynamics of the propagating gravitational field let us now turn to a simpler problem and inquire about the physical phenomena of the non-propagating, near field sector. The simplest electrodynamics problem for fixed external charges is that of a point charge with the solution of the Coulomb force. The analogous problem, the gravitational field created by a static point mass is the Schwarzschild solution. We use the notation $x^0 = ct \rightarrow t$ in this section.

A. Metric

The symmetry of the corresponding space-time is time independence and rotational invariance. The rotational invariance requires that (i) the space-time can be foliated by a family of two-dimensional surfaces, $\Sigma(t, r)$ and (ii) that any pair of points of a hyper-surface there is a spatial rotation bringing one point into the other. The time independence assures that there is a time-like unit vector field $n(x)$, $n^2(x) = 1$, generating an infinitesimal transformation of the space-time,

$x^\mu \rightarrow x^\mu + \epsilon n^\mu(x)$, which leaves the geometry, the metric tensor in particular, invariant. The spatial rotations generate displacement orthogonal to the time direction defined by the vector field $n^\mu(x)$ therefore the most general static, rotational invariant metric is of the form

$$ds^2 = f(\mathbf{x})dt^2 - \sum_{j,k=1}^3 x^j h_{jk}(\mathbf{x})x^k \quad (216)$$

where $h_{jk}(\mathbf{x})$ is a three-dimensional, rotational invariant metric. Let us chose the radial coordinate

$$r = \sqrt{\frac{A}{4\pi}} \quad (217)$$

where A is the time independent area of the surface $\Sigma(t, r)$ and parameterize this surface by the polar angle parameters θ and ϕ , giving

$$ds^2 = f(r)dt^2 - h(r)dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (218)$$

The non-vanishing Christoffel symbols of this metric are

$$\begin{aligned} \Gamma^t_{rt} &= \frac{f'}{2f}, & \Gamma^r_{tt} &= \frac{f'}{2h} \\ \Gamma^r_{rr} &= \frac{h'}{2h}, & \Gamma^r_{\theta\theta} &= -\frac{r}{h}, & \Gamma^r_{\phi\phi} &= -\frac{r \sin^2 \theta}{h}, & \Gamma^\theta_{r\theta} &= \Gamma^\phi_{r\phi} = \frac{1}{r} \\ \Gamma^\theta_{\phi\phi} &= -\sin \theta \cos \theta, & \Gamma^\phi_{\theta\phi} &= \cot \theta \end{aligned} \quad (219)$$

and the other non-vanishing components can be obtained by exchanging the covariant indices. The Ricci tensor is diagonal,

$$\begin{aligned} R_{tt} &= -\frac{f''}{2h} + \frac{f'}{4h} \left(\frac{f'}{f} + \frac{h'}{h} \right) - \frac{f'}{rh} \\ R_{rr} &= \frac{f''}{2f} - \frac{f'}{4f} \left(\frac{f'}{f} + \frac{h'}{h} \right) - \frac{h'}{rh} \\ R_{\theta\theta} &= -1 + \frac{r}{2h} \left(\frac{f'}{f} - \frac{h'}{h} \right) + \frac{1}{h} \\ R_{\phi\phi} &= R_{\theta\theta} \sin^2 \theta. \end{aligned} \quad (220)$$

The vacuum Einstein equation for $r \neq 0$ where $R = 0$ are

$$R_{tt} = R_{rr} = R_{\theta\theta} = 0. \quad (221)$$

Since

$$0 = \frac{R_{tt}}{f} + \frac{R_{rr}}{h} = -\frac{1}{rh} \left(\frac{f'}{f} + \frac{h'}{h} \right) \quad (222)$$

we have

$$\frac{f'}{f} + \frac{h'}{h} = 0 \quad (223)$$

requiring

$$hf = A, \quad (224)$$

A being a constant. The condition that the metric approaches the flat one for $r \rightarrow \infty$ gives $A = 1$ and find for the last equation in (221)

$$0 = rf' + f - 1 \quad (225)$$

or

$$\frac{drf}{dr} = 1. \quad (226)$$

The solution is

$$f = 1 + \frac{B}{r} \quad (227)$$

where B is a constant. The parameterization $B = -2GM/c^2$ yields the metric

$$ds^2 = \left(1 - \frac{r_s}{r}\right) dt^2 - \frac{dr^2}{1 - \frac{r_s}{r}} - r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (228)$$

where

$$r_s = \frac{2GM}{c^2} \quad (229)$$

is the Schwarzschild-radius (with $c \neq 1$ restored). It is advantageous to remove the dimension of the coordinates by the intrinsic scale r_s , $t \rightarrow tr_s$ and $r \rightarrow rr_s$ and write the dimensionless invariant length square as

$$ds^2 = \left(1 - \frac{1}{r}\right) dt^2 - \frac{dr^2}{1 - \frac{1}{r}} - r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (230)$$

Few remarks are in order at this point.

1. The gravitational field is weak for $r \gg 1$ and Newton's gravitational law applies approximately according to Eq. (175).
2. The proper time of a stationary observer which stays at a given spatial coordinate, $d\tau = \sqrt{1 - \frac{1}{r}} dt < dt$ shows that freely falling clocks slow down, suffer a red-shift as $r \rightarrow 1$ from above.

3. A stationary observer's four-velocity is $u_t = \frac{1}{\sqrt{1-\frac{1}{r}}}$, $u_r = u_\theta = u_\phi = 0$. Its four-acceleration,

$$a^\mu = u^\nu D_\nu u^\mu = u^\nu \partial_\nu a^\mu + \Gamma^\mu_{\rho\nu} u^\rho u^\nu = \frac{\Gamma^\mu_{tt}}{1 - \frac{1}{r}}, \quad (231)$$

has a single non-vanishing component, $a^r = \frac{1}{2r^2} = \phi'(r)$ where $\phi(r) = -\frac{1}{2r}$ is the Newtonian potential according to Eq. (175), canceling the gravitational force, in agreement with the Equivalence Principle. But the regularity of the acceleration is misleading because the physical, gauge invariant acceleration is obtained by multiplying it by $\sqrt{g_{rr}}$,

$$\sqrt{-a^\mu g_{\mu\nu} a^\nu} = \frac{1}{2r^2 \sqrt{1 - \frac{1}{r}}}, \quad (232)$$

which diverges at $r = 1$: It is more difficult to remain stationary as the observer approaches $r = 1$ from above.

4. The metric shows two singularities, one at the Schwarzschild radius

$$r_s \approx \begin{cases} 2.8 \frac{M}{M_{sun}} \text{ km} \\ 2.4 \frac{M}{M_{proton}} \cdot 10^{-52} \text{ cm} \end{cases} \quad (233)$$

and another at $r = 0$. The former turns out to be a singularity of this coordinate system because the curvature tensor remain regular and can be eliminated by means of appropriately defined coordinates. The latter is a true singularity.

5. Despite the absence of a singularity something dramatic happens at the Schwarzschild radius. The light cone corresponds to the infinitesimal changes

$$\frac{dt}{dr} = \pm \frac{1}{1 - \frac{1}{r}}, \quad (234)$$

the light cones have space-dependent orientation which becomes singular at $r = 1$, the causal structure changes discontinuously at the Schwarzschild radius. Another pathology is seen by considering the motion of a massive particle where $ds^2 > 0$. This inequality is consistent with constant r , the particle can be at rest compared to the Schwarzschild radius if $r > 1$. But this is not possible anymore for $r < 1$ where the role of the time t as a coordinate with positive unit vector is taken over the the radius r and thereby is forced to change during the motion.

6. The solution requires that the mass be concentrated at $r = 0$. For any realistic, non-singular mass distribution the solution is more complicated in the space region with non-vanishing




FIG. 2: The light-cone structure of time-like geodesics and an inward (a) and outward (b) massive particle world line.

mass density. The Schwarzschild-radius is naked and visible only for mass distributions which are vanishing for $r \geq 1$. The numerical values given in Eq. (233) suggest that the Schwarzschild-radius might be found experimentally in astrophysics rather than in particle physics.

7. The solution remains the same when time independence is not assumed at the beginning, namely the spherically symmetric solutions of the vacuum Einstein equation are static (Kirchhoff's theorem). This holds in the Newtonian theory in an obvious manner since the mass can be concentrated at the origin for spherically symmetrical field. This theorem excludes the spherically symmetric s -waves from gravitational radiation field.

B. Geodesics

Let us consider the motion of a massive particle in the Schwarzschild geometry where the Lagrangian

$$L = -mc\sqrt{\dot{x}^\mu g_{\mu\nu}(x)\dot{x}^\nu} \quad (235)$$

can be written as

$$L = -mc\sqrt{\left(1 - \frac{1}{r}\right)\dot{t}^2 - \frac{\dot{r}^2}{1 - \frac{1}{r}} - r^2(\dot{\theta}^2 + \sin^2\theta\dot{\phi}^2)}. \quad (236)$$

The motion is always planar, equation of motion for θ ,

$$r^2 \sin\theta \cos\theta \dot{\phi}^2 = \frac{d}{ds} r^2 \dot{\theta} \quad (237)$$

is satisfied by $\theta(s) = \pi/2$, the case considered hereafter. One usually solves the non-relativistic problem by exploiting the energy conservation. Since the temporal component of the relativistic equation of motion is the energy conservation such a starting point corresponds to the use of the equation $\dot{x}^2 = \kappa$, giving

$$\left(1 - \frac{1}{r}\right) \dot{t}^2 - \frac{\dot{r}^2}{1 - \frac{1}{r}} - r^2 \dot{\phi}^2 = \kappa. \quad (238)$$

We set $\kappa = 1$ in this calculation of the orbit of a massive object. The coordinates t and ϕ are cyclic therefore the corresponding momentum is conserved,

$$\begin{aligned} -\frac{1}{mc} \frac{\partial L}{\partial \dot{t}} &= \left(1 - \frac{1}{r}\right) \dot{t} = E, \\ \frac{1}{mc} \frac{\partial L}{\partial \dot{\phi}} &= r^2 \dot{\phi} = L, \end{aligned} \quad (239)$$

and (238) gives

$$\frac{E^2 - \dot{r}^2}{1 - \frac{1}{r}} - \frac{L^2}{r^2} = \kappa \quad (240)$$

what we write

$$\dot{r}^2 + V(r) = E^2 \quad (241)$$

in terms of the effective potential

$$V(r) = \left(1 - \frac{1}{r}\right) \left(\kappa + \frac{L^2}{r^2}\right). \quad (242)$$

The contribution proportional to $\kappa = 1$ is the Newtonian effective potential and the remaining $\mathcal{O}(r^{-3})$ piece represents relativistic effects. The motion with $|E| < 1$ is bounded, $0 < r < r_{max}$, because $V(0) = -\infty$ and $V(\infty) = 1$. The extrema of the potential satisfy the equation

$$\frac{dV(r)}{d\frac{1}{r}} = -3\frac{L^2}{r^2} + 2\frac{L^2}{r} - 1 = 0. \quad (243)$$

The potential is monotonous when $L < \sqrt{3}$ and the particle falls into the center. For $L > \sqrt{3}$ the potential displays a local maximum at are

$$r_{max} = \frac{3}{1 + \sqrt{1 - \frac{3}{L^2}}}, \quad (244)$$

and a local minimum at

$$r_{min} = \frac{3}{1 - \sqrt{1 - \frac{3}{L^2}}}, \quad (245)$$




FIG. 3: The effective potential, (242), as the function of r/r_s for $L^2/r_s^2 = 1.5, \sqrt{3}, 1.85, 2.0$ in increasing order.

cf. Fig. 3 and there are stable orbits in certain range of E .

The presence of the $\mathcal{O}(r^{-3})$ relativistic term in the effective potential (242) violates Kepler's law for planetary motion, in particular it induces a perihelion motion. This brought the first decisive victory for General relativity when Einstein could reproduce the perihelion motion of Mercury, known since in 1859.

The Euler-Lagrange equation of the Lagrangian of a massive particle, (163), gives a geodesics. One expects that the world line of a light particle approaches the motion of a massless particle when the mass tends to zero. This can be easily established in an obvious manner for photons in the geometrical limit by means of Fermat's principle. By assuming here that massless particles follow null-geodesics the previous consideration remains valid with $\kappa = 0$. Photons with $\kappa = 0$ do not feel the Newtonian gravitational potential as expected but their orbital angular momentum is coupled to gravitation. This is not surprising since the affine connection term in the covariant derivative couples the polarization of the electromagnetic radiation to gravity. The polarization follows the change of the direction of propagation and couples the orbital angular momentum to gravity. The deflection of light around the Sun has been observed first in 1919 and was the second major support of General Relativity.

C. Space-like hyper-surfaces

The Schwarzschild geometry is static and its non-trivial features are captured by the constant time hyper-surfaces, a one-dimensional manifold of curved three-dimensional spaces, parametrized by t . Each three-dimensional hyper-surface can be embedded into a four dimensional Euclidean space. To simplify matters we consider the two-dimensional $\theta = \pi/2$ section of the hyper-surfaces, obeying the metric

$$-ds^2 = \frac{dr^2}{1 - \frac{1}{r}} + r^2 d\phi^2. \quad (246)$$

This surface can easily be embedded into a three-dimensional Euclidean space by means of the cylindrical coordinates (z, r, ϕ) . The surface $z = z(r)$ in the Euclidean three-space space of metric

$$-ds^2 = dz^2 + dr^2 + r^2 d\phi^2 \quad (247)$$

has the invariant length

$$-ds^2 = [1 + z'^2(r)] dr^2 + r^2 d\phi^2, \quad (248)$$

where $z'(r) = \frac{dz(r)}{dr}$. The comparison with (246) gives

$$1 + z'^2(r) = \frac{1}{1 - \frac{1}{r}}, \quad (249)$$

and

$$z(r) = \int_1^r \frac{dr'}{\sqrt{r' - 1}} = 2\sqrt{r - 1}, \quad (250)$$

defined for $r > 1$ only.

D. Around the Schwarzschild-horizon

The causal structure of space-time is determined by the local light cones because any signal or interaction can propagate on their surface or within them. The light cones, given by Eq. (234) become narrow and narrow as the Schwarzschild-radius is approached from above, as shown in Fig. 2. This indicates that the free fall motion, seen by a stationary observer slows down as the horizon is approached. This can be understood as a manifestation of the red-shift, mentioned in point 2. of Section VI A. The role of the radial and the time coordinate of the metric is exchanged for a time-like geodesic as it traverses $r = 1$ and the light-cones are oriented horizontally. The Kruskal-Szekeres coordinate system, introduced below shows that such an orientation of the light-cones makes that no physical object of signal can crosses $r = 1$ from below. Therefore the sphere $r = 1$ is a horizon which can be traversed inside only and appears for the outside observers as a black hole.

The metric (230) has a coordinate singularity only at the Schwarzschild-radius because the non-vanishing components of the curvature tensor

$$R_{t\theta t\theta} = R_{t\phi t\phi} = -R_{r\theta r\theta} = -R_{r\phi r\phi} = \frac{1}{2}R_{\theta\phi\theta\phi} = -\frac{1}{2}R_{trtr} = \frac{1}{r^3} \quad (251)$$

make the eigenvalues, the invariant content of the curvature tensor, regular. Therefore a point particle experiences nothing irregular or special when crossing the horizon. But an object which




FIG. 4: (a): The geometrical origin of the parametrization (254). (b): The radius r as the function of the proper time, s , for the free fall with $L = 0$ from $r = 2$.

is extended in the radial direction suffers strong tidal forces at the horizon. This is because a small, finite separation, Δr , corresponds to diverging invariant length, $\sqrt{-\Delta s^2} = \Delta r / \sqrt{1 - 1/r}$, as $r \rightarrow 1$.

1. Falling through the horizon

It is instructive to follow the radial free fall of a point particle through the horizon. The equation of motion of a massive particle for $L = 0$ is

$$\dot{r}^2 = \frac{1}{r} + E^2 - 1. \quad (252)$$

Let us suppose that the motion starts with vanishing velocity at $r = r_0 > 1$ and write

$$ds = \frac{dr}{\sqrt{\frac{1}{r} - \frac{1}{r_0}}}. \quad (253)$$

Such a $r(s)$ function can easily be obtained in a parametrized form,

$$\begin{aligned} r &= \frac{r_0}{2}(1 + \cos \eta), \\ s &= \frac{r_0^{3/2}}{2}(\eta + \sin \eta), \end{aligned} \quad (254)$$

known from the description of the motion of a point on a circle, rolling with constant speed. Nothing special happens at $r = 1$ and the total proper time of falling into the center at $\eta = \pi$ is $s = \frac{\pi}{2}r_0^{3/2}$, cf. Fig. 4.

2. Stretching the horizon

We have seen so far, that the free fall, followed by its proper time shows no particular singularity at the horizon. But what does a stationary observer finds which uses the Schwarzschild coordinates




FIG. 5: The tortoise coordinate, r^* as function of r .

t and r ? The Schwarzschild time can be found from the first equation of (239),

$$\dot{r} = \frac{dr}{dt} \dot{t} = \frac{dr}{dt} \frac{E}{1 - \frac{1}{r}}, \quad (255)$$

indicating a singularity for such an observer when the horizon is crossed. Such a singularity might be avoided by the use of the coordinate r^* satisfying

$$dr^* = \frac{dr}{1 - \frac{1}{r}}, \quad (256)$$

because the singular factor is absorbed in the new coordinate and our equation for the radius now reads as

$$\dot{r} = \frac{dr^*}{dt} E. \quad (257)$$

The solution of Eq. (256),

$$r^* = r + \ln|r - 1|, \quad (258)$$

replaced into the equation of motion, (252), gives

$$E^2 \left[1 - \left(\frac{dr^*}{dt} \right)^2 \right] = 1 - \frac{1}{r}. \quad (259)$$

If $r \rightarrow 1$ from above then $r^* \rightarrow -\infty$,

$$\frac{dr^*}{dt} \rightarrow -1 \quad (260)$$

and $t \sim -r^* \rightarrow \infty$, it takes infinitely long time to fall through the Schwarzschild-radius. It is easy to understand that this is the result of the apparent singularity in the Schwarzschild metric, $g_{tt} \rightarrow 0$, $g_{rr} \rightarrow \infty$ with $g_{tt}g_{rr} = 1$ as $r \rightarrow 1$.

The radius r^* , defined by Eq. (258) is called tortoise coordinate. We are told that Achilles could not pass a tortoise because each time he reached the point where the tortoise was before it was already ahead. The singularity of the Schwarzschild metric, the factor $1/(1 - 1/r)$ multiplying

dr^2 in the expression of the invariant length (230), signals that the scale of the radius should be refined, r should be allowed to decrease beyond zero to describe the free fall through the horizon. We have $r^* \sim r$ and $r^* \rightarrow -\infty$ as $r \rightarrow 1$, shown qualitatively in Fig. 5, stretching out conveniently the approach of the horizon from either side.

3. Crossing the horizon

The Schwarzschild time diverges on both sides of the horizon as indicated in Fig. 2. To resolve the traverse the horizon we need better suited coordinates. Instead of the coordinates t and r one may use $u = t - r$ and $v = t + r$ in Minkowski flat space-time, labeling the out- and in-going light rays, respectively. The expression of the invariant length is $ds^2 = dt^2 - dr^2 = dudv$, showing clearly that the new coordinates correspond to light cones because $ds^2 = 0$ for $du = 0$ or $dv = 0$. The Eddington-Finkelstein coordinates are based on such a change of coordinates in the Schwarzschild space-time except that r is replaced by the tortoise coordinate to resolve the approach the horizon,

$$\tilde{u} = t - r^*, \quad \tilde{v} = t + r^*, \quad (261)$$

yielding the expression

$$\begin{aligned} ds^2 &= \left(1 - \frac{1}{r}\right) d\tilde{v}^2 - 2d\tilde{v}dr - r^2(d\theta^2 + \sin^2\theta d\phi^2) \\ &= \left(1 - \frac{1}{r}\right) d\tilde{u}^2 + 2d\tilde{u}dr - r^2(d\theta^2 + \sin^2\theta d\phi^2). \end{aligned} \quad (262)$$

The null-geodesics satisfy the equations

$$\frac{d\tilde{v}}{dr} = \begin{cases} 0 & in, \\ \frac{2}{1-\frac{1}{r}} & out, \end{cases}, \quad \frac{d\tilde{u}}{dr} = \begin{cases} 0 & out, \\ -\frac{2}{1-\frac{1}{r}} & in, \end{cases} \quad (263)$$

where the direction of the geodesics, in or out, is determined by the help of Fig. 5. The in- and out-going null-geodesics form horizontal lines on the (r, \tilde{v}) and (r, \tilde{u}) planes, respectively. The other, non-trivially oriented null-geodesics are $\tilde{v}_{out} = -\tilde{v}_{in} = 2r^*$, depicted qualitatively in Figs. 6.

4. Reduplication of space-time

Though the Eddington-Finkelstein coordinates describes the in- and out-going geodesics we need different representations, (\tilde{v}, r) and (\tilde{u}, r) , respectively. To have common coordinate system for both set of geodesics we drop r and use \tilde{u} and \tilde{v} and the expression

$$ds^2 = \left(1 - \frac{1}{r}\right) d\tilde{u}d\tilde{v} - r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (264)$$




FIG. 6: The in- and out- going null-geodesics on the planes (a): (r, \tilde{u}) and (b): (r, \tilde{v}) as function of r/r_r . The dashed line stands at the horizon and the arrows indicate the direction of the time.

for time and radius dependent part of the invariant length. This is indeed a simple metric but shows a singularity at the horizon. But this singularity is now less troublesome, a simple rescaling of the coordinates,

$$\begin{aligned} (I) \quad u' &= -e^{-\frac{\tilde{u}}{2}} = -\sqrt{r-1}e^{\frac{r-t}{2}}, & v' &= e^{\frac{\tilde{v}}{2}} = \sqrt{r-1}e^{\frac{r+t}{2}}, \\ (II) \quad u' &= e^{-\frac{\tilde{u}}{2}} = \sqrt{1-re}e^{\frac{r-t}{2}}, & v' &= e^{\frac{\tilde{v}}{2}} = \sqrt{1-re}e^{\frac{r+t}{2}}, \end{aligned} \quad (265)$$

where the transformations (I) and (II) apply outside and inside of the Schwarzschild sphere, respectively, remove the singularity since the metric in terms of the new dimensionless coordinate,

$$ds^2 = \frac{4}{r}e^{-r} du' dv' - r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (266)$$

is regular at $r = 1$. The coordinates u' and v' correspond to null-directions, it is more natural to use the time- and space-like coordinates,

$$\rho = \frac{v' - u'}{2}, \quad \tau = \frac{v' + u'}{2}, \quad (267)$$

given by

$$\begin{aligned} (I) \quad \rho &= \sqrt{r-1}e^{\frac{r}{2}} \cosh \frac{t}{2}, & \tau &= \sqrt{r-1}e^{\frac{r}{2}} \sinh \frac{t}{2}, \\ (II) \quad \rho &= \sqrt{1-re}e^{\frac{r}{2}} \sinh \frac{t}{2}, & \tau &= \sqrt{1-re}e^{\frac{r}{2}} \cosh \frac{t}{2}, \end{aligned} \quad (268)$$

yielding the metric

$$ds^2 = \frac{4}{r}e^{-r}(d\tau^2 - d\rho^2) - r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (269)$$

The final problem to overcome is that this transformation covers a part of the space-time only because it gives $v' > 0$. The regions with $v' < 0$ can be obtained by using

$$\begin{aligned} (III) \quad u' &= e^{-\frac{\tilde{u}}{2}} = \sqrt{r-1}e^{\frac{r-t}{2}}, & v' &= -e^{\frac{\tilde{v}}{2}} = -\sqrt{r-1}e^{\frac{r+t}{2}}, \\ (IV) \quad u' &= -e^{-\frac{\tilde{u}}{2}} = \sqrt{1-re}e^{\frac{r-t}{2}}, & v' &= -e^{\frac{\tilde{v}}{2}} = -\sqrt{1-re}e^{\frac{r+t}{2}}, \end{aligned} \quad (270)$$




FIG. 7: The mapping of two Schwarzschild space-time onto the Kruskal-Szekeres geometry. The constant time and radius curves are radial or horizontally opening hyperbolic curves, respectively, such as the dotted lines and $t = -\infty \rightarrow \rho = -\tau$, $t = 0 \rightarrow \tau = 0$, $t = \infty \rightarrow \rho = \tau$. The outside and inside of the horizon $r = r_s \rightarrow \rho = \pm\tau$, is mapped into (I)-(III) and (II)-(IV), respectively and the Schwarzschild geometry, (I)-(II) is reduplicated into (III)-(IV).

instead of (265), yielding

$$\begin{aligned}
 (III) \quad \rho &= -\sqrt{r-1}e^{\frac{r}{2}} \cosh \frac{t}{2}, & \tau &= -\sqrt{r-1}e^{\frac{r}{2}} \sinh \frac{r}{2}, \\
 (IV) \quad \rho &= -\sqrt{1-re^{\frac{r}{2}}} \sinh \frac{t}{2}, & \tau &= -\sqrt{1-re^{\frac{r}{2}}} \cosh \frac{t}{2}.
 \end{aligned} \tag{271}$$

The transformation, given by eqs. (268) and (271) defines the Kruskal-Szekeres coordinates, satisfying

$$(r-1)e^r = \rho^2 - \tau^2 \quad (I), (II), (III) \text{ and } (IV), \quad t = \begin{cases} 2\operatorname{arcth} \frac{\tau}{\rho} & (I) \text{ and } (III), \\ 2\operatorname{arcth} \frac{\rho}{\tau} & (II) \text{ and } (IV), \end{cases} \tag{272}$$

cf. Fig. 7.

The matching of the two space-times in the outer region can be demonstrated by the Einstein-Rose bridge, the extension of the embedding, described in Section VI C. One chooses an Euclidean three-space, parametrized by the coordinates (z, u, ϕ) and equipped by the metric

$$-ds^2 = dz^2 + d\rho^2 + r^2 d\phi^2. \tag{273}$$

The surface $z = z(\rho)$ has the induced metric

$$-ds^2 = [1 + z'^2(\rho)]d\rho^2 + r^2 d\phi^2, \tag{274}$$

which is to be matched to (269),

$$-ds^2 = \frac{4}{r}e^{-r} d\rho^2 + r^2 d\phi^2. \tag{275}$$




FIG. 8: The Einstein-Rosen bridge, connecting the two space-times in the $(z, \rho \cos \phi, \rho \sin \phi)$ coordinate system. The regions (I) and (III) belong to $z > 0$ and $z < 0$, respectively. A strip of the surface $\rho = \infty$ is indicated by the dotted lines, they form the horizontal part of a bended plane. The bending represents the analytical continuation between the regions (I) and (III). It is supposed to be at infinitely far, leaving the throat, indicated by the circles, ϕ -independent, rotation invariant. The dashed line follows the path $0 < \rho < \infty$ in (I) and (III).

Hence the equation

$$1 + z'^2(\rho) = \frac{4}{r} e^{-r} \quad (276)$$

follows. The solution, as a function of the Schwarzschild radius r ,

$$z(r) = \int_1^r dr' \frac{d\rho}{dr} \left(\frac{4}{r'} e^{-r'} - 1 \right), \quad (277)$$

is sketched qualitatively in Fig. 8. It remains a challenge to understand the effects of such a dramatic reduplication of the Universe at each point particle.

5. Causal structure

The causal structure can easily be recognized in the Kruskal-Szekeres space-time because τ is a time-like coordinate everywhere and the light-cones preserve their direction, contrary to the Schwarzschild parametrization, displayed in Fig. 2. The equal time lines indicate that the Schwarzschild time t runs in opposite directions in the two space-times, (I)-(II) and (III)-(IV). One can see immediately that the two outer regions, (I) and (III) are causally disconnected.

Though time and space coordinates appear in equal footing in relativity, the time has a distinguished role, it parametrizes the motion and has an orientation, encoded by the time arrow. The radial ($L = 0$) null geodesic, line cones, are parallel to the coordinate axes u' and v' and two forward oriented light cones of a particle, slightly before and after crossing the Schwarzschild




FIG. 9: The future light cones on the Kruskal-Szekeres space-time, indicated by the dotted lines running parallel with the u' and v' axes. The fat solid line represents the singular point $r = 0$.

radius are indicated in Fig. 9. One can see that before and after the crossing the ingoing light rays fall into the singularity but the outward oriented light rays stay outside of the horizon or fall later into the center, respectively. The massive particle world lines remain within the future light cones, therefore no particle, either massive or massless, is able to emerge from the region $r < 1$ to $r > 1$: The Schwarzschild-sphere is impenetrable from inside. Note that the horizon is free of singularities and its one-way oriented, irreversible passage results from an appropriate rearrangement of the future light cones without strong forces.

VII. HOMOGENEOUS AND ISOTROPIC COSMOLOGY

The assumption that we do not occupy any special location in the Universe suggests that there should be a coordinate system in which the matter distribution in the Universe appears homogeneous on large enough distance scales. Furthermore, the absence of a preferred direction suggests isotropy, as well. The astrophysical observations supports these assumptions with an astonishing precision.

A gauge or coordinate system independent way of stating spatial homogeneity is to impose the existence of a one dimensional family of hyper-surfaces, $\Sigma(t)$ which (i) foliate the space-time, ie. any space-time point corresponds to one and only one hyper-surface and (ii) for any pair of points $p, q \in \Sigma(t)$ there is an isometry, a scalar product preserving mapping of the space-time which sends p into q .

Isotropy states that for any pair of space-like unit vectors, $v(p)$ and $v'(p)$ at a given space-time point p there is an isometry of the space-time which rotates $v(p)$ into $v'(p)$. The Universe appears

isotropic in good approximation.

We introduce the concept of standard observers which find the distant galaxies at rest. Their world lines provides a time-like congruence, system of time-like curves which fill up the space-time.

It is worthwhile mentioning two remarks at this point. (i) The tangent vectors of the time-like congruence are orthogonal to the tangent vectors of the space-like hyper-surfaces of the homogeneity assumption at each space-time point. In fact, otherwise one could construct a preferred spatial direction which contradicts isotropy. (ii) The space-time metric g induces a three-dimensional metric h on the hyper-surfaces. The isometry which maps a point of a hyper-surface $\Sigma(t)$ into another one according to the assumption of homogeneity is clearly an isometry of this three-dimensional induced metric, too. Furthermore, according to the previous remark this three-dimensional geometry has no preferred directions.

A. Maximally symmetric spaces

The homogeneous and isotropic three-space at a given time appears as symmetric as possible. To make this concept more precise let us consider geometries with maximal number of symmetries, Killing vectors.

The reparametrization $x^\mu \rightarrow x'^\mu = x^\mu - w^\mu(x)$ is a symmetry of the metric tensor if the deformation $w(x)$ is a Killing field, satisfying the Killing equation, (116). The parallel transport along an infinitesimal rectangle gives the equation

$$D_\mu D_\nu w_\rho - D_\nu D_\mu w_\rho = -R^\lambda_{\rho\mu\nu} w_\lambda. \quad (278)$$

There are two similar equations, obtained by cyclic permutation of the indices,

$$\begin{aligned} D_\nu D_\rho w_\mu - D_\rho D_\nu w_\mu &= -R^\lambda_{\mu\nu\rho} w_\lambda, \\ D_\rho D_\mu w_\nu - D_\mu D_\rho w_\nu &= -R^\lambda_{\nu\rho\mu} w_\lambda. \end{aligned} \quad (279)$$

the sum of these three equation can be written for a Killing field as

$$D_\mu D_\nu w_\rho - D_\nu D_\mu w_\rho + D_\rho D_\mu w_\nu = 0, \quad (280)$$

due to cyclic symmetry (93). The expression (278) of the parallel transport allows us to rewrite this equation in the form

$$D_\rho D_\mu w_\nu = R^\lambda_{\rho\mu\nu} w_\lambda, \quad (281)$$

stating that second (covariant) derivative of the Killing field can be expressed in terms of the Killing field itself, w . The x -dependence is therefore given by the value of the Killing field and its first derivative at a given point, $w^\mu(x_0)$, $D_\nu w^\mu(x_0)$. We need yet another property of the vector fields, a set of fields $\{w_n^\mu(x)\}$ is called independent if the vanishing of the linear superposition, made up by constant coefficient,

$$\sum_n c_n w_n^\mu(x) = 0 \quad (282)$$

implies $c_n = 0$.

We have d independent vectors $w^\mu(x_0)$ and $d(d-1)/2$ independent anti-symmetric tensors $D_\nu w_\mu(x_0) - D_\mu w_\nu(x_0)$ at each point in d -dimensions hence the maximally symmetric space has $d + d(d-1)/2 = d(d+1)/2$ independent Killing vectors. For instance the d -dimensional Euclidean space has d translational and $d(d-1)/2$ rotational symmetries. Furthermore, homogeneous and isotropic spaces have maximal symmetry.

B. Robertson-Walker metric

Let us consider now the three-dimensional curvature tensor, $\tilde{R}_{k\ell m}^j$, more precisely the tensor

$$\tilde{R}_{\ell m}^{jk} = \tilde{R}_{n\ell m}^j h^{kn} \quad (283)$$

which can be thought as a transformation of second order antisymmetric contravariant tensors,

$$T^{\ell k} = -T^{k\ell} \rightarrow \tilde{R}_{mn}^{k\ell} T^{mn}. \quad (284)$$

The matrix \tilde{R} which acts on the tensor space is symmetric according to the second line in Eqs. (110) therefore it can be diagonalized. Isotropy of the tree-space, the absence of preferred direction requires that the matrix of this map be degenerate,

$$\tilde{R}_{\ell m}^{jk} = K(\delta_\ell^j \delta_m^k - \delta_\ell^k \delta_m^j). \quad (285)$$

The degenerate eigenvalue K is related to the scalar curvature,

$$\tilde{R} = k|^{(d)}R| = Kd(d-1), \quad (286)$$

with $k = -1, 0, 1$ in d spatial dimensions. The corresponding Ricci tensor is

$$\tilde{R}_m^k = \tilde{R}_{jm}^{jk} = \frac{\tilde{R}}{d} \delta_m^k = K(d-1)\delta_m^k \quad (287)$$

The spatial homogeneity makes \tilde{R} constant within each spatial hyper-surface $\Sigma(t)$.

We shall now construct the metric on a spatial hyper-surface with homogeneous curvature. Spaces with positive curvature, $k = 1$, can be obtained by embedding into R^4 ,

$$x^2 + y^2 + z^2 + w^2 = a^2, \quad (288)$$

yielding

$$0 = xdx + ydy + zdz + wdw \quad (289)$$

and the induced metric

$$\begin{aligned} ds^2 &= dx^2 + dy^2 + dz^2 + dw^2 \\ &= dx^2 + dy^2 + dz^2 + \frac{(xdx + ydy + zdz)^2}{a^2 - x^2 - y^2 - z^2} \end{aligned} \quad (290)$$

written in polar coordinates as

$$\begin{aligned} ds^2 &= dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2) + \frac{r^2 dr^2}{a^2 - r^2} \\ &= \frac{dr^2}{1 - \frac{r^2}{a^2}} + r^2(d\theta^2 + \sin^2 \theta d\phi^2). \end{aligned} \quad (291)$$

For negative curvature the embedding is given by

$$x^2 + y^2 + z^2 - w^2 = -a^2, \quad (292)$$

which yields

$$0 = xdx + ydy + zdz - wdw \quad (293)$$

and the induced metric is

$$\begin{aligned} ds^2 &= dx^2 + dy^2 + dz^2 - dw^2 \\ &= dx^2 + dy^2 + dz^2 - \frac{(xdx + ydy + zdz)^2}{a^2 + x^2 + y^2 + z^2}. \end{aligned} \quad (294)$$

We find in polar coordinates

$$\begin{aligned} ds^2 &= dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2) - \frac{r^2 dr^2}{a^2 + r^2} \\ &= \frac{dr^2}{1 + \frac{r^2}{a^2}} + r^2(d\theta^2 + \sin^2 \theta d\phi^2). \end{aligned} \quad (295)$$

The spatially homogeneous and isotropic space-time provides time coordinate axis orthogonal to the space directions, therefore the four dimensional metric is of Robertson-Walker,

$$ds^2 = d\tau^2 - a^2(\tau) \left[\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2 \theta d\phi^2) \right] \quad (296)$$

where τ is the proper time measured by clocks in rest in the space-like hyper-surface, the coordinate r is made dimensionless by means of the scale factor $r \rightarrow a(u)r$ which is an arbitrary constant for flat space and $k = \text{sign}K$. The three-space is of finite volume for positive curvature, $k = 1$ and infinite for $k = 0, -1$.

One may write the metric as

$$ds^2 = d\tau^2 - a^2(\tau) [d\chi^2 + h^2(\chi)(d\theta^2 + \sin^2 \theta d\phi^2)] \quad (297)$$

with

$$r = h(\chi) = \begin{cases} \sin \chi & k = 1 \\ \chi & k = 0 \\ \sinh \chi & k = -1 \end{cases} \quad (298)$$

which justifies the use of the combination

$$r_{ph} = a(\tau)h(\chi) \quad (299)$$

as a cosmic distance parameter. Yet another useful form of the metric is

$$ds^2 = a^2(\eta) [d\eta^2 - d\chi^2 - h^2(\chi)(d\theta^2 + \sin^2 \theta d\phi^2)] \quad (300)$$

where

$$\eta = \int \frac{d\tau}{a(\tau)}. \quad (301)$$

The metric tensor

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 \\ 0 & -a^2(t)\tilde{g}_{ij} \end{pmatrix}, \quad \tilde{g}_{ij} = \begin{pmatrix} \frac{1}{1-kr^2} & 0 & 0 \\ 0 & r^2 & 0 \\ 0 & 0 & r^2 \sin^2 \theta \end{pmatrix}, \quad (302)$$

$\mu, \nu = (\tau, r, \theta, \phi)$,

$$\sqrt{-g} = a^3(\tau) \frac{r^2 \sin \theta}{\sqrt{1-kr^2}} \quad (303)$$

yields the Christoffel symbols with two or more indices τ vanishing and the further non-vanishing components are

$$\Gamma_{ij}^\tau = \dot{a}a\tilde{g}_{ij}, \quad \Gamma_{\tau j}^i = \frac{\dot{a}}{a}\tilde{g}_j^i, \quad \Gamma_{jk}^i = \tilde{\Gamma}_{jk}^i = \frac{1}{2}\tilde{g}^{i\ell}(\partial_j\tilde{g}_{k\ell} + \partial_k\tilde{g}_{\ell j} - \partial_\ell\tilde{g}_{jk}) \quad (304)$$

with $\dot{a} = \frac{da}{d\tau}$. The Ricci tensor of the metric tensor \tilde{g} is proportional to \tilde{g} itself and the proportionality constant turns out to be after some calculation $k(d-1)$ for a d -dimensional homogeneous and isotropic space. Thus we have

$$\tilde{R}_{jm} = 2k\tilde{g}_{jm} \quad (305)$$

for the three-space and

$$R_{\mu\nu} = \begin{pmatrix} R_{00} & 0 \\ 0 & \tilde{R}_{jk} + \tilde{g}_{jk}(a\ddot{a} + 2\dot{a}^2) \end{pmatrix} = \begin{pmatrix} -3\frac{\ddot{a}}{a} & 0 \\ 0 & \tilde{g}_{jk}(a\ddot{a} + 2\dot{a}^2 + 2k) \end{pmatrix} \quad (306)$$

for the four dimensional space-time. The scalar curvature is

$$R = R_{00} - \frac{1}{a^2}\tilde{g}^{jk}R_{jk} = -6\left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2}\right). \quad (307)$$

C. Equation of motion

In order to find the Einstein equation we approximate the matter, averaged over long distances, as an ideal fluid which is rest in the cosmic coordinate system of the Robertson-Walker metric, ie. the energy-momentum tensor is given by

$$T^{\mu\nu} = (\rho c^2 + p)u^\mu u^\nu - pg^{\mu\nu}, \quad (308)$$

where u^μ is the four-velocity of the matter, the unit tangent vector field of the time-like congruence. We have $T^\mu_\mu = \rho c^2 - 3p = 0$ for scale invariant case like EM radiation, $p = 0$ for matter at rest, like cosmic dust and $T^{\mu\nu} \sim g^{\mu\nu}$, $p = -\rho c^2$ in the vacuum. Notice that $\rho c^2 + 3p \geq 0$ in each case. The metric (296) is based on the time coordinate τ therefore $u^\mu = (1, 0, 0, 0)$.

The first two terms in the divergence of a tensor $A^{\mu\nu}$

$$D_\nu A^{\mu\nu} = \partial_\nu A^{\mu\nu} + \Gamma^\nu_{\rho\nu} A^{\mu\rho} + \Gamma^\mu_{\rho\nu} A^{\rho\nu} \quad (309)$$

looks as the covariant divergence of a four-vector which can be written in a simpler manner according to (123),

$$D_\nu A^{\mu\nu} = \frac{1}{\sqrt{-g}}\partial_\nu(\sqrt{-g}A^{\mu\nu}) + \Gamma^\mu_{\rho\nu}A^{\rho\nu}. \quad (310)$$

Thus the expression (308) leads to the energy-momentum conservation law

$$0 = -\partial_\nu pg^{\mu\nu} + \frac{1}{\sqrt{-g}}\partial_\nu[\sqrt{-g}(\rho c^2 + p)u^\mu u^\nu] + \Gamma^\mu_{\rho\nu}(\rho c^2 + p)u^\rho u^\nu \quad (311)$$

where the metric admissibility, $Dg = 0$, was used, too. The rest frame condition, $u^\mu = (1, 0, 0, 0)$, renders the spatial components, $\mu = 1, 2, 3$ of this equation trivial and the temporal part $\mu = 0$ reads as

$$a^3 \dot{p} = \frac{d}{d\tau} [a^3 (\rho c^2 + p)], \quad (312)$$

giving

$$0 = \frac{d}{d\tau} (a^3 \rho c^2), \quad (313)$$

$\rho \sim 1/a^3$ for dust. In the case of radiation we write

$$0 = \frac{4}{3} \frac{d}{d\tau} (a^3 \rho c^2) - \frac{1}{3} a^3 \dot{\rho} c^2 = \frac{1}{a} \frac{d}{d\tau} (a^4 \rho c^2), \quad (314)$$

resulting in $\rho \sim 1/a^4$. The density drops faster in the latter case during the expansion of the universe (growing a) than for dust. Though the radiation represents a negligible component in the actual universe, it was dominant in an earlier phase.

The Einstein equations read finally as

$$\begin{aligned} R_{00} - \frac{1}{2} R - \Lambda &= 3 \frac{\dot{a}^2 + k}{a^2} - \Lambda \\ &= 8\pi G T_{\tau\tau} = 8\pi G \rho c^2 \end{aligned} \quad (315)$$

for the components 00 and

$$\begin{aligned} \frac{1}{a^2 \tilde{g}_{rr}} \left[R_{rr} + \frac{a^2 \tilde{g}_{rr}}{2} (R + 2\Lambda) \right] &= \frac{\ddot{a}}{a} + 2 \frac{\dot{a}^2}{a^2} + 2 \frac{k}{a^2} - 3 \left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right) + \Lambda \\ &= -2 \frac{\ddot{a}}{a} - \frac{\dot{a}^2}{a^2} - \frac{k}{a^2} + \Lambda \\ &= \frac{8\pi G}{a^2 \tilde{g}_{rr}} T_{rr} = 8\pi G p \end{aligned} \quad (316)$$

for rr .

We can express the acceleration \ddot{a} by making forming a suitable linear superposition of these two equations,

$$\frac{\ddot{a}}{a} = \frac{\Lambda}{3} - \frac{4}{3} \pi G (3p + \rho c^2). \quad (317)$$

The cosmological constant introduces a pressure in the absence of matter and leads to violation of the Newtonian gravitational law in the slow motion, weak gravitational field limit. We shall set $\Lambda = 0$ in the rest of the discussion for simplicity.

The first remark is that there is no static solution, $\ddot{a} < 0$, for $3p + \rho c^2 > 0$. The rate of change of spatial distances,

$$v = \frac{d\ell}{d\tau} = \frac{\ell}{a}\dot{a} = H\ell, \quad (318)$$

where

$$H = \frac{\dot{a}}{a} \quad (319)$$

called Hubble-constant, though its value has slow time dependence on astrophysical time scale. It was supposed to be around 600km/s/Mpc according as proposed by Hubble in the '30s, its present value is around 70km/s/Mpc. Note that $v > c$ for large enough separation. This is not in contradiction with special relativity which considers velocities at the same space-time point but leads to the appearance of horizons as we shall see later.

The universe is expanding at the present, $\dot{a} > 0$ but in view of $\ddot{a} < 0$ the expansion rate must have been faster in the past. By assuming constant expansion rate $T = 1/H$ time ago the universe would have been point-like. Due to the slowing expansion rate the big bang, the zero-size universe must have occurred less time before and the inverse Hubble-constant gives only an order of magnitude estimate of the lifetime of the universe. The zero size signals a singularity in the time evolution which prevents us to inquire about the state of the universe before the big bang. The so called singularity theorems of general relativity assures that the singularity at the big bang is present even without assuming homogeneity and isotropy.

For the flat or open universe, $k = 0$ or $k = -1$, respectively $\dot{a} \neq 0$ according to Eq. (315) which can be written as

$$\dot{a}^2 = \frac{8\pi G}{3}a^2\rho c^2 - k \quad (320)$$

and the expansion continues forever. In fact, $\rho c^2 = \mathcal{O}(a^{-3})$ or $\rho c^2 = \mathcal{O}(a^{-4})$ for dust or radiation dominated universe, $\rho c^2 a^2 \rightarrow 0$ as $\tau \rightarrow \infty$ and \dot{a} approaches zero from above. For closed universe, $k = 1$, the matter contribution to Eq. (320) decreases compared to k during the expansion and there is a maximal value of a , $a \leq a_0$. But the maximal value can not be approached asymptotically because \ddot{a} does not tend to zero according to Eq. (317) but instead a big crunch occurs at some finite time where $a = 0$ is reached and the universe ceases to exist.

The 00 component of the Einstein equation (315) for $\Lambda = 0$ shows that the universe is closed or open if $\rho > \rho_c$ or $\rho < \rho_c$, respectively where

$$\rho_c c^2 = \frac{3H^2}{8\pi G}. \quad (321)$$

The actual observational and theoretical background suggests that the cosmological constant Λ actually plays an important role in determining the age of the universe, in particular the choice $\rho_{matter} \approx 0.27\rho_c$, $\rho_\Lambda \approx 0.73\rho_c$, $\rho_{matter} + \rho_\Lambda \approx \rho_c$ is preferred.

D. Frequency shifts

The precision spectroscopic measurement of the electromagnetic radiation from stars is an easy way of collecting information about distant astrophysical objects. Three different mechanisms are known to change the observed value of the characteristic frequency of a physical system. The first is observed in flat space-time when the source of a wave and its observer moves with respect to each other. The second and the the third mechanisms are for sources and observers at rest but in the presence of a static or time-dependent gravitational field, respectively.

1. Let us consider a monochromatic plane wave with wave vector $k^\mu = (\omega_0/c, \mathbf{k})$ with $k^2 = m^2c^2/\hbar^2$ whose source is moving with velocity \mathbf{v} with respect to an observer in flat space-time. The time component of the wave vector in the observer's reference frame is

$$\omega = \frac{\omega_0 - \mathbf{v}\mathbf{k}}{\sqrt{1 - \frac{v^2}{c^2}}}. \quad (322)$$

By means of the relation $|\mathbf{k}| = \sqrt{\frac{\omega_0^2}{c^2} - k^2}$ we write

$$\mathbf{v}\mathbf{k} = v \frac{\omega_0}{c} \sqrt{1 - \frac{m^2c^4}{\hbar^2\omega_0^2}} \cos \theta \quad (323)$$

and find

$$\omega = \omega_0 \frac{1 - \frac{v}{c} \sqrt{1 - \frac{m^2c^4}{\hbar^2\omega_0^2}} \cos \theta}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (324)$$

This is the relativistic Doppler effect, its non-relativistic analogy for a simple wave propagating with speed \tilde{c} with $v, \tilde{c} \ll c$ is

$$\omega = \omega_0 \left(1 - \frac{v}{\tilde{c}} \cos \theta\right). \quad (325)$$

The Doppler shift is particularly useful in the observation of the broadening of spectral lines of light arriving from stars because it allows us to estimate the random velocity of the atoms or molecules and the temperature on the surface of the star.

2. Time independent gravitational field changes the frequency, being proportional with the energy. Let us consider two electron-positron pair at rest far from a star of mass M . One pair

annihilates into a photon and the other starts to fall freely towards the star. Let us suppose that at distance r from the star the second pair annihilates into a photon. The photon arising from the first pair, at large distance from the star, is of frequency $\omega_1(\infty) = E_k(\infty)/\hbar = 2m_0c^2/\hbar$. The photon which is created at distance r by the second pair is of the frequency $\omega_2(r) = E_k(r)/\hbar$ where

$$E_k = 2 \left(m_0c^2 + \frac{Gm_0M}{r} \right) = 2m_0c^2 \left(1 + \frac{GM}{rc^2} \right) \quad (326)$$

is the kinetic energy of the pair at the instance of the annihilation. Let us suppose that we send the photon arising from the annihilation of the first pair at the location where the annihilation of the second pair took place by means of static mirrors which leave the photon frequency unchanged and denote its frequency $\omega_1(r)$ there. Energy conservation requires $\omega_1(r) = \omega_2(r)$ since otherwise one can extract energy from static gravitational field by annihilation of electron-positron pairs at certain distance from the star, sending the photons arising from the process at a different distance and recombining them into electron-positron pair again. Thus the frequency of the photon changes as

$$\omega(r) = \omega(\infty) \left(1 + \frac{GM}{rc^2} \right) \quad (327)$$

due to the presence of the gravitational field. The relative change,

$$\frac{\Delta\omega}{\omega} = \frac{\omega(r) - \omega(\infty)}{\omega(\infty)} = \frac{GM}{rc^2} \quad (328)$$

is approximately $2 \cdot 10^{-6}$ at the surface of the Sun but becomes comparable to one on the surface of a neutron star and can be used to estimate the total mass.

3. Time dependent gravitational field leads to the change of frequency in a trivial manner because the time of emission of a photon is different than the time of its absorption. Let us suppose that a light signal is emitted at the point $p_0 = (ct_e, \mathbf{r})$ which is received at $p = (ct_o, \mathbf{0})$. The propagation is along a null-geodesic,

$$cdt = a(ct)d\chi, \quad (329)$$

yielding

$$\int_{t_e}^{t_o} \frac{cdt}{a(ct)} = \chi. \quad (330)$$

The source emits $dn = dt_e\omega_e$ periods during the time interval dt_e of the emission with frequency ω_e . It is the same number of periods, $dn = dt_o\omega_o$ which is observed in time dt_o and frequency ω_o . Thus we have red shift parameter, the relative change of the wavelength $\lambda = 2\pi c/\omega$,

$$z = \frac{\lambda_o - \lambda_e}{\lambda_e} = \frac{dt_o}{dt_e} - 1 = \frac{a(ct_o)}{a(ct_e)} - 1 \quad (331)$$

where the difference of Eq. (330) and its analogy written for the times $t_e \rightarrow t_e + dt_e$, $t_o \rightarrow t_o + dt_o$ was used in the last equation. By assuming that the scale factor $a(\tau)$ changes slowly in time we find

$$\begin{aligned} z &= \frac{a_o}{a_o + (ct_e - ct_o)\dot{a}_o + \frac{1}{2}(ct_e - ct_o)^2\ddot{a}_o + \dots} - 1 \\ &= \frac{\dot{a}_o}{a_o}(ct_o - ct_e) + \left[\left(\frac{\dot{a}_o}{a_o} \right)^2 - \frac{\ddot{a}_o}{2a_o} \right] (ct_o - ct_e)^2 + \dots \end{aligned} \quad (332)$$

where $a_o = a(ct_o)$. The deceleration parameter,

$$q = -\frac{a\ddot{a}}{\dot{a}^2} \quad (333)$$

measures the rate of change of the Hubble-constant in time and its experimental value is about 1. Hubble's law, the linearity of the red shift in the distance is obtained by

$$z \approx \frac{\dot{a}_o}{a_o}(ct_o - ct_e) \approx \frac{\dot{a}_o}{a_o}\ell = H\ell. \quad (334)$$

As a simple application let us now consider the absolute luminosity \mathcal{L} of a galaxy and the measured flux (energy per unit time and unit area) \mathcal{F} . The quantity

$$d_L = \sqrt{\frac{\mathcal{L}}{4\pi\mathcal{F}}} \quad (335)$$

can be interpreted as the luminosity distance of the galaxy. But due to the expansion of the universe the distance of the galaxy at the time of the emission of the observed signal, r_e was different than r_o , the distance at the time of observation. Does the distance d_L agree one of them? Not. Energy conservation requires

$$\mathcal{F} = \frac{\mathcal{L}}{4\pi a^2(ct_o)r_e^2} = \frac{\mathcal{L}}{4\pi a^2(ct_o)r_e^2(1+z)^2}, \quad (336)$$

where the factor $dA/4\pi a^2(ct_o)r_e^2$ is the solid angle of a surface dA observed, one factor of $1+z$ arises from the red shift of the observed photons and the other from the ratio of the time intervals dt_e/dt_o and we find

$$d_L = a(ct_o)r_e(1+z). \quad (337)$$

E. Particle horizon

Can we receive signals from any part of the universe? It is true that the universe was smaller in the past but the speed of light appeared smaller, too. This is the kind of dilemma which led to Eq.

(335) and requires more careful consideration. We now use the form (300) of the metric and look for the spatial region from which signals can be received in the case of flat geometry, $k = 0$. The metric is now conformally flat, we can replace the overall conformal factor $a^2(\eta)$ by one from the point of view of this question and the problem is reduced to flat Minkowski geometry. Obviously if the integral

$$\int_{\epsilon}^{\tau} \frac{d\tau'}{a(\tau')} \quad (338)$$

remains finite when $\epsilon \rightarrow 0$ then we can not receive signals from the whole universe and there is a particle horizon. The short time solution of the Einstein equations give $a(\tau) = \mathcal{O}(\tau^{2/3})$ for $\tau \approx 0$ even for a dust dominated universe and the particle horizon appears. For $k = -1$ the curvature term becomes negligible for short time and the same result is recovered. The situation is more complicated for a closed universe and it turns out that the horizon disappears when the universe reaches its maximal size for a dust domination but remains present for all times in the radiation dominated case.

The presence of horizons raises serious problems for the description of the evolution of the universe. The reason is that the difference between regions of the early universe which are separated by horizon can not relax during the evolution and any inhomogeneity which appeared at such scale in the very early universe should be observable today. The cosmic microwave background which is supposed to originate from a rather early period of the universe is found to be homogeneous and isotropic to such a large extent which is not possible to understand unless one assumes that either the universe is created in an unusually homogeneous state or the Robertson-Walker metric is not valid for the early phase. The inflationary universe model which involves a rapid expansion in the early phase leads to an enlargement of the horizon and offers a possibility to reconcile the entropy estimates of the early universe with the present homogeneity of the microwave background radiation.

But the latest, more sensitive measurements of the anisotropy and inhomogeneities of the microwave background radiation lead to another serious problem, these fluctuations are apparently too large to be explained by the inflationary standard model. Further problems which point to the need of modifying the Einstein equation at the scale of the planetary system or astrophysical length come from the Pioneer anomaly, an inexplicable deviation of the Pioneer sonde from the orbits predicted by Einstein's general relativity and the more and more precisely determined slowing down of the expansion of the universe. This latter phenomenon might be explained by assuming that the majority of the matter in our universe is participating in gravitational interaction only

and pulls back the expanding, visible matter. But such a rescue operation of the Einstein equation seems to be out of proportion and the slight modification of the Einstein-Hilbert action appears to be a more economical and better justifiable way of establishing consistency.

F. Evolution of the universe

We shall briefly review the salient feature of the evolutionary big bang model. It is clearly unreasonable to expect that classical physics, in particularly general relativity as we know it today is capable to trace the evolution just from the beginning, from $t = 0$, $a = 0$. The basic physical constants can be put together to form a length scale,

$$\ell_P = \sqrt{\frac{G\hbar}{c^3}} \approx 10^{-33} \text{ cm}, \quad (339)$$

called Planck length. At distances smaller than this scale the gravitational interaction should be stronger than the quantum effects and the question of quantizing gravity, neither necessary according to the experiments nor understood today, becomes a central issue. In general, one should not extrapolate the physical laws across such a vast regime of scale what separate the length scale 10^{-16} cm, the space resolution of the the experiments today, from the Planck length because the basic structure of the physical laws changed in the last century several time by the discovery of new interactions or particles as the observational length scale was reduced from the mm scale to the proton size, 10^{-13} cm. It is a real surprise that the structure of quantum mechanics found to be valid in this range when all other laws in physics proved to be limited to longer length scales.

The first question one might raise after leaving the **Planck-era** ($t \approx 10^{-43}$ s, $T \approx 10^{31}$ K, $\rho \approx 10^{92}$ gm/cm³) is whether thermodynamical equilibrium was reached by the universe. The inflation, the rapid expansion of the universe can provide such a thermalization by the enlargement of the horizons. The background microwave radiation is the indication that thermal equilibrium was reached by a hot universe. According to Eqs. (313) and (314) the early universe was dominated by radiation.

Another interesting issue of this period is the generation of matter-anti matter domains. Our galaxy tend to have matter rather than anti-matter, the baryon number (+1 for a nucleon and -1 for an anti-nucleon) its total baryon number, a conserved quantity, is positive. If the universe was created by positive net baryon number then this is not surprising. But it is more natural to imagine that the universe started its existence with vanishing conserved charges and the matter

excess observed around us is compensated by an anti-matter excess somewhere else in the universe. What was the mechanism which created these matter-anti matter domains? There is no generally accepted and satisfactory answer.

The following milestones should be mentioned for the subsequent evolution:

1. At the beginning after the Planck-era the non-gravitational interactions are supposed to be unified by the **Grand Unified Model**. At about $t \approx 10^{-34}$ s, $T \approx 10^{26}$ K, $\rho \approx 10^{68}$ gm/cm³ the symmetry of the Grand Unified Model is broken spontaneously, the strong and the electroweak interactions separate into Quantum Chromodynamics and the unified electroweak theory, respectively and the matter-anti matter islands are supposed to be formed.
2. At about $t \approx 10^{-12}$ s, $T \approx 10^{15}$ K, $\rho \approx 10^{31}$ gm/cm³ the symmetry of the **unified electroweak theory** is broken spontaneously and electromagnetic and weak forces separate.
3. At $t \approx 10^{-6}$ s, $T \approx 10^{13}$ K, $\rho \approx 10^{17}$ gm/cm³ quarks which were freely propagating became confined and the present day **hadrons are formed**.
4. At $t \approx 1$ s, $T \approx 10^{10}$ K, $\rho \approx 10^8$ gm/cm³ the universe consists of mainly neutrinos, photons, electrons, protons and neutrons and their anti-particles were in thermal equilibrium up to now but the interaction with neutrinos becomes weak to maintain equilibrium from now on. As a result **neutrinos decouple** and follow a passive red shift in the rest of the time. Soon after the **proton-neutron conversion is frozen** out ($T = 10^{10}$ K corresponds approximately to 1MeV, the neutron-proton mass difference), too.
5. At $t \approx 4$ s, $T \approx 5 \cdot 10^9$ K, $\rho \approx 3 \cdot 10^4$ gm/cm³ the **electron-positron equilibrium is lost** because $T = T \approx 5 \cdot 10^9$ K belongs to 0.5MeV, the mass of the electron. The annihilation process eliminates positrons and heats up the photons slightly.
6. At $t \approx 3$ min, $T \approx 10^9$ K, $\rho \approx 10$ gm/cm³ the thermal energy reaches the nuclear binding scale and **nucleosynthesis** starts by producing ⁴He nuclei. The strong Coulomb barrier and the lack of other stable elements with $Z < 8$ leave helium the only nuclei produced in mass, until all neutrons left over after step 4. are bound into helium. At the end of the helium dominated era which lasts few minutes 25% of the mass is in the form of ⁴He, the rest is essentially distributed over ²H, ³He and ⁷Li. The reproduction of the fraction taken in the form of ⁴He is a convincing success of the hot big bang nucleosynthesis model.

7. At $t \approx 4 \cdot 10^5$ year, $T \approx 5 \cdot 10^4$ K, $\rho \approx 10^{-16}$ gm/cm³ the **matter reaches equilibrium with radiation** and starts to dominate.
8. At $t \approx 10^6$ year, $T \approx 5 \cdot 2 \cdot 10^3$ K, $\rho \approx 10^{-19}$ gm/cm³ the thermal energy reaches the ionization energy of the hydrogen atom and we enter in the recombination era when stable, **neutral atoms**, starting with the hydrogen are formed. Most of the universe becoming electrically neutral the photons decouple. Their subsequent cooling leads to the actual temperature 2.7K, observed with high accuracy in the **cosmic microwave background** in the last decades. This is the second decisive victory of the hot big bang model for the universe.

The decoupling of matter and radiation signals the end of the **quantum-driven evolution** era. The resulting loss of radiation pressure leads to gravitational instabilities for masses $M > 10^5 M_{sun}$ where $M_{sun} \approx 2 \cdot 10^{33}$ g is the solar mass and **galaxies** start to be formed. About $t \approx 10^3 - 10^7$ years hadronic matter started to dominate the total energy of the universe whose present age is currently believed to be $14 \cdot 10^9$ years.

Appendix A: Classical Field theory

The goal of this Appendix is a brief introduction to variational principle and conservation laws in classical field theory. The reason to go beyond the usual, differential equation based definition of the dynamics is have equations of motion which preserve their form under arbitrary transformation of the space-time coordinates.

1. Variational principle

Field theory is a dynamical system containing degrees of freedom, denoted by $\phi(\mathbf{x})$, at each space point \mathbf{x} . The coordinate $\phi(\mathbf{x})$ can be a single real number (real scalar field) or consist n -components (n -component field). Our goal is to provide an equation satisfied by the trajectory $\phi_{cl}(t, \mathbf{x})$. The index cl is supposed to remind us that this trajectory is the solution of a classical equation of motion. The problem of identifying $\phi_{cl}(t, \mathbf{x})$ will be outlined in three steps.

a. *Single point on the real axis*

Problem: identification of a point on the real axis, $x_{cl} \in \mathbb{R}$, in a manner which is independent of the reparametrization of the real axis.

Solution: Find a function with vanishing derivative at x_{cl} only:

$$\frac{df(x)}{dx} \Big|_{x=x_{cl}} = 0 \quad (\text{A1})$$

To check the reparametrization invariance of this equation we introduce new coordinate y by the function $x = x(y)$ and find

$$\frac{df(x(y))}{dy} \Big|_{y=y_{cl}} = \underbrace{\frac{df(x)}{dx} \Big|_{x=x_{cl}}}_{0} \frac{dx(y)}{dy} \Big|_{y=y_{cl}} = 0 \quad (\text{A2})$$

Variational principle: There is simple way of rewriting Eq. (A1). Let us perform an infinitesimal variation of the coordinate $x \rightarrow x + \delta x$, and write

$$\begin{aligned} f(x_{cl} + \delta x) &= f(x_{cl}) + \delta f(x_{cl}) \\ &= f(x_{cl}) + \delta x \underbrace{f'(x_{cl})}_0 + \frac{\delta x^2}{2} f''(x_{cl}) + \mathcal{O}(\delta x^3) \end{aligned} \quad (\text{A3})$$

The variation principle, equivalent of Eq. (A1) is

$$\delta f(x_{cl}) = \mathcal{O}(\delta x^2), \quad (\text{A4})$$

stating that x_{cl} is characterized by the property that an infinitesimal variation around it, $x_{cl} \rightarrow x_{cl} + \delta x$, induces an $\mathcal{O}(\delta x^2)$ change in the value of $f(x_{cl})$.

b. *Non-relativistic point particle*

Problem: identification of a trajectory in a coordinate choice independent manner.

Variational principle: Let us identify a trajectory $x_{cl}(t)$ by specifying the coordinate at the initial and final time, $x_{cl}(t_i) = x_i$, $x_{cl}(t_f) = x_f$ (by assuming that the equation of motion is of second order in time derivatives) and consider a variation of the trajectory $x(t)$: $x(t) \rightarrow x(t) + \delta x(t)$ which

leaves the initial and final conditions invariant (ie. does not modify the solution). Our function $f(x)$ of the previous section becomes a functional, called action

$$S[x(\cdot)] = \int_{t_i}^{t_f} dt L(x(t), \dot{x}(t)) \quad (\text{A5})$$

involving the Lagrangian $L(x(t), \dot{x}(t))$. (The symbol $x(\cdot)$ in the argument of the action functional is supposed to remind us that the variable of the functional is a function. It is better to put a dot in the place of the independent variable of the function $x(t)$ otherwise $\left. \begin{array}{l} \text{]} \\ \text{]]} \end{array} \right\}$

The coordinate not appearing in the Lagrangian in an explicit manner is called cyclic coordinate,

$$\frac{\partial L(x, \dot{x})}{\partial x_{cycl}} = 0. \quad (\text{A13})$$

For each cyclic coordinate there is a conserved quantity because the generalized momentum of a cyclic coordinate, p_{cycl} is conserved according to Eqs. (A11) and (A13).

c. Scalar field

Problem: identification of the equation of motion for an n -component field, $\phi_a(x)$, $a = 1, \dots, n$.

(Notation: $x = (t, \mathbf{x})$.)

Variational principle: let us consider a variation of the trajectory $\phi(x)$:

$$\phi(x) \rightarrow \phi(x) + \delta\phi(x), \quad \delta\phi(t_i, \mathbf{x}) = \delta\phi(t_f, \mathbf{x}) = 0. \quad (\text{A14})$$

The variation of the action

$$S[\phi(\cdot)] = \int_V dt d^3x L(\phi, \partial\phi) \quad (\text{A15})$$

is

$$\begin{aligned} \delta S &= \int_V dt d^3x \left(\frac{\partial L(\phi, \partial\phi)}{\partial \phi_a} \delta\phi_a + \frac{\partial L(\phi, \partial\phi)}{\partial \partial_\mu \phi_a} \delta \partial_\mu \phi_a \right) + \mathcal{O}(\delta^2 \phi) \\ &= \int_V dt d^3x \left(\frac{\partial L(\phi, \partial\phi)}{\partial \phi_a} \delta\phi_a + \frac{\partial L(\phi, \partial\phi)}{\partial \partial_\mu \phi_a} \partial_\mu \delta\phi_a \right) + \mathcal{O}(\delta^2 \phi) \\ &= \int_{\partial V} ds^\mu \delta\phi_a \frac{\partial L(\phi, \partial\phi)}{\partial \partial_\mu \phi_a} + \int_V dt d^3x \delta\phi_a \left(\frac{\partial L(\phi, \partial\phi)}{\partial \phi_a} - \partial_\mu \frac{\partial L(\phi, \partial\phi)}{\partial \partial_\mu \phi_a} \right) + \mathcal{O}(\delta^2 \phi) \end{aligned} \quad (\text{A16})$$

The first term for $\mu = 0$,

$$\int_{\partial V} ds^0 \delta\phi_a \frac{\partial L(\phi, \partial\phi)}{\partial \partial_0 \phi_a} = \int_{t=t_f} d^3x \underbrace{\delta\phi_a}_{0} \frac{\partial L(\phi, \partial\phi)}{\partial \partial_0 \phi_a} - \int_{t=t_i} d^3x \underbrace{\delta\phi_a}_{0} \frac{\partial L(\phi, \partial\phi)}{\partial \partial_0 \phi_a} = 0 \quad (\text{A17})$$

is vanishing because there is no variation at the initial and final time. When $\mu = j$ then

$$\int_{\partial V} ds^j \delta\phi_a \frac{\partial L(\phi, \partial\phi)}{\partial \partial_j \phi_a} = \int_{x_j=\infty} ds^j \delta\phi_a \underbrace{\frac{\partial L(\phi, \partial\phi)}{\partial \partial_j \phi_a}}_0 - \int_{x_j=-\infty} ds^j \delta\phi_a \underbrace{\frac{\partial L(\phi, \partial\phi)}{\partial \partial_j \phi_a}}_0 = 0 \quad (\text{A18})$$

and it is still vanishing because we are interested in the dynamics of localized systems and the interactions are supposed to be short ranged. Therefore, $\phi = 0$ at the spatial infinities and the Lagrangian is vanishing. The suppression of the second term gives the Euler-Lagrange equation

$$\frac{\partial L(\phi, \partial\phi)}{\partial \phi_a} - \partial_\mu \frac{\partial L(\phi, \partial\phi)}{\partial \partial_\mu \phi_a} = 0. \quad (\text{A19})$$

Examples:

1. Free scalar particle:

$$L = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{m^2}{2} \phi^2 \quad \Longrightarrow \quad (\square + m^2) \phi(x) = 0 \quad (\text{A20})$$

2. Self interacting scalar particle:

$$L = \frac{1}{2} (\partial \phi)^2 - \frac{m^2}{2} \phi^2 - \mathfrak{V}(\phi) \quad \Longrightarrow \quad (\partial_\mu \partial^\mu + m^2) \phi = -\mathfrak{V}'(\phi) \quad (\text{A21})$$

2. Noether theorem

The reparametrization invariance of the Euler-Lagrange equation shows that there is a conserved current for each continuous symmetry.

Symmetry: A transformation of the space-time coordinates $x^\mu \rightarrow x'^\mu$, and the field $\phi_a(x) \rightarrow \phi'_a(x)$ preserves the equation of motion. Since the equation of motion is obtained by varying the action, the action should be preserved by the symmetry transformations. A slight generalization is that the action can in fact be changed by a surface term which does not influence its variation, the equation of motion at finite space-time points. Therefore, the symmetry transformations satisfy the condition

$$L(\phi, \partial \phi) \rightarrow L(\phi', \partial' \phi') + \partial'_\mu \Lambda^\mu \quad (\text{A22})$$

with a certain vector function $\Lambda^\mu(x')$.

Continuous symmetry: There are infinitesimal symmetry transformations, in an arbitrary small neighborhood of the identity, $x^\mu \rightarrow x^\mu + \delta x^\mu$, $\phi_a(x) \rightarrow \phi_a(x) + \delta \phi_a(x)$. Examples: Rotations, translations in the space-time, and $\phi(x) \rightarrow e^{i\alpha} \phi(x)$ for a complex field.

Conserved current: $\partial_\mu j^\mu = 0$, conserved charge: $Q(t)$:

$$\partial_0 Q(t) = \partial_0 \int_V d^3 x j^0 = - \int_V d^3 x \partial_\nu j^\nu = - \int_{\partial V} ds \cdot \mathbf{j} \quad (\text{A23})$$

It is useful to distinguish external and internal spaces, corresponding to the space-time and the values of the field variable. Eg.

$$\phi_a(x) : \underbrace{\mathbb{R}^4}_{\text{external space}} \rightarrow \underbrace{\mathbb{R}^m}_{\text{internal space}} . \quad (\text{A24})$$

Internal and external symmetry transformations act on the internal or external space, respectively.

a. *Point particle*

The main points of the construction of the Noether current for internal symmetries can be best understood in the framework of a particle.

To find the analogue of the internal symmetries let us consider a point particle with the continuous symmetry $\mathbf{x} \rightarrow \mathbf{x} + \epsilon \mathbf{f}(\mathbf{x})$ for infinitesimal ϵ ,

$$L(\mathbf{x}, \dot{\mathbf{x}}) = L(\mathbf{x} + \epsilon \mathbf{f}(\mathbf{x}), \dot{\mathbf{x}} + \epsilon(\dot{\mathbf{x}} \cdot \partial) \mathbf{f}(\mathbf{x})) + \mathcal{O}(\epsilon^2). \quad (\text{A25})$$

Let us introduce a new, time dependent coordinates, $\mathbf{y}(t) = \mathbf{y}(\mathbf{x}(t))$, based on the solution of the equation of motion, $\mathbf{x}_{cl}(t)$, in such a manner that one of them will be $y^1(t) = \epsilon(t)$, where $\mathbf{x}(t) = \mathbf{x}_{cl}(t) + \epsilon(t) \mathbf{f}(\mathbf{x}_{cl}(t))$. There will be $n - 1$ other new coordinates, y^ℓ , $\ell = 2, \dots, n$ whose actual form is not interesting for us. The Lagrangian in terms of the new coordinates is defined by $L(\mathbf{y}, \dot{\mathbf{y}}) = L(\mathbf{y}(\mathbf{x}), \dot{\mathbf{y}}(\mathbf{x}))$. The ϵ -dependent part assumes the form

$$L(\epsilon, \dot{\epsilon}) = L(\mathbf{x}_{cl} + \epsilon \mathbf{f}(\mathbf{x}_{cl}), \dot{\mathbf{x}}_{cl} + \epsilon(\dot{\mathbf{x}}_{cl} \cdot \partial) \mathbf{f}(\mathbf{x}_{cl}) + \dot{\epsilon} \mathbf{f}(\mathbf{x}_{cl})) + \mathcal{O}(\epsilon^2). \quad (\text{A26})$$

What is the equation of motion of this Lagrangian? Since the solution is $\epsilon(t) = 0$ it is sufficient to retain the $\mathcal{O}(\epsilon)$ contributions in the Lagrangian only,

$$L(\epsilon, \dot{\epsilon}) \rightarrow L^{(1)}(\epsilon, \dot{\epsilon}) = \epsilon \frac{\partial L(\mathbf{x}_{cl}, \dot{\mathbf{x}}_{cl})}{\partial \mathbf{x}} \cdot \mathbf{f}(\mathbf{x}_{cl}) + \frac{\partial L(\mathbf{x}_{cl}, \dot{\mathbf{x}}_{cl})}{\partial \dot{\mathbf{x}}} [\epsilon(\dot{\mathbf{x}}_{cl} \cdot \partial) \mathbf{f}(\mathbf{x}_{cl}) + \dot{\epsilon} \mathbf{f}(\mathbf{x}_{cl})] \quad (\text{A27})$$

up to an ϵ -independent constant. The corresponding Euler-Lagrange equation is

$$\frac{\partial L^{(1)}(\epsilon, \dot{\epsilon})}{\partial \epsilon} - \frac{d}{dt} \frac{\partial L^{(1)}(\epsilon, \dot{\epsilon})}{\partial \dot{\epsilon}} = 0. \quad (\text{A28})$$

(this is the point where the formal invariance of the equation of motion under nonlinear, time dependent transformations of the coordinates is used). According to Eq. (A25) ϵ is a cyclic coordinate,

$$\frac{\partial L(\epsilon, \dot{\epsilon})}{\partial \epsilon} = 0 \quad (\text{A29})$$

and its generalized momentum,

$$p_\epsilon = \frac{\partial L(\epsilon, \dot{\epsilon})}{\partial \dot{\epsilon}} \quad (\text{A30})$$

is conserved.

The external space transformation corresponds to the shift of the time, $t \rightarrow t + \epsilon$ which induces $x(t) \rightarrow x(t - \epsilon) = x(t) - \epsilon \dot{x}(t)$ for infinitesimal ϵ . This is a symmetry as long as the Hamiltonian (and

the Lagrangian) does not contain explicitly the time. In fact, the action changes by a boundary contribution only which can be seen by expanding the Lagrangian in time around $t - \epsilon$,

$$\int_{t_i}^{t_f} dt L(x(t), \dot{x}(t)) = \int_{t_i}^{t_f} dt \left[L(x(t - \epsilon), \dot{x}(t - \epsilon)) + \epsilon \frac{dL(x(t), \dot{x}(t))}{dt} \right] + \mathcal{O}(\epsilon^2) \quad (\text{A31})$$

and as a result the variational equation of motion remains unchanged. But the continuation of the argument is slightly different from the case of internal symmetry. We consider ϵ as a time dependent function which generates a transformation of the coordinate, $x(t) \rightarrow x(t - \epsilon(t)) = x(t) - \epsilon(t)\dot{x}(t) + \mathcal{O}(\epsilon^2)$. The Lagrangian of $\epsilon(t)$ as new coordinate for $x(t) = x_{cl}(t)$ is

$$\begin{aligned} L^{(1)}(\epsilon, \dot{\epsilon}) &= L(x_{cl}(t - \epsilon), \dot{x}_{cl}(t - \epsilon)) - L(x_{cl}(t), \dot{x}_{cl}(t)) \\ &= -\epsilon \dot{x}_{cl} \frac{\partial L(x_{cl}, \dot{x}_{cl})}{\partial x} - \frac{d\epsilon \dot{x}_{cl}}{dt} \frac{\partial L(x_{cl}, \dot{x}_{cl})}{\partial \dot{x}} + \mathcal{O}(\epsilon^2) \\ &= \underbrace{-\epsilon \dot{x}_{cl} \frac{\partial L(x_{cl}, \dot{x}_{cl})}{\partial x} - \epsilon \ddot{x}_{cl} \frac{\partial L(x_{cl}, \dot{x}_{cl})}{\partial \dot{x}} - \dot{\epsilon} \dot{x}_{cl} \frac{\partial L(x_{cl}, \dot{x}_{cl})}{\partial \dot{x}}}_{-\epsilon \frac{dL(x_{cl}, \dot{x}_{cl})}{dt}} + \mathcal{O}(\epsilon^2) \\ &= -\epsilon \left[\frac{dL(x_{cl}, \dot{x}_{cl})}{dt} - \frac{d}{dt} \left(\frac{\partial L(x_{cl}, \dot{x}_{cl})}{\partial \dot{x}} \dot{x}_{cl} \right) \right] - \frac{d}{dt} \left(\frac{\partial L(x_{cl}, \dot{x}_{cl})}{\partial \dot{x}_{cl}} \epsilon \dot{x}_{cl} \right) + \mathcal{O}(\epsilon^2) \end{aligned} \quad (\text{A32})$$

up to an ϵ -independent constant. Its Euler-Lagrange equation (A28) assures the conservation of the energy,

$$H = \frac{\partial L(x, \dot{x})}{\partial \dot{x}} \dot{x} - L(x, \dot{x}). \quad (\text{A33})$$

b. Internal symmetries

An internal symmetry transformation of field theory acts on the internal space only. We shall consider linearly realized internal symmetries for simplicity where

$$\delta x^\mu = 0, \quad \delta \phi_a(x) = \epsilon \underbrace{\tau_{ab}}_{generator} \phi_b(x). \quad (\text{A34})$$

This transformation is a symmetry,

$$L(\phi, \partial\phi) = L(\phi + \epsilon\tau\phi, \partial\phi + \epsilon\tau\partial\phi) + \mathcal{O}(\epsilon^2). \quad (\text{A35})$$

Let us introduce new "coordinates", ie. new field variable, $\Phi(\phi)$, in such a manner that $\Phi^1(x) = \epsilon(x)$ where $\phi(x) = \phi_{cl}(x) + \epsilon(x)\tau\phi_{cl}(x)$, $\phi_{cl}(x)$ being the solution of the equations of movement. The linearized Lagrangian for $\epsilon(x)$ is

$$\begin{aligned} L(\epsilon, \partial\epsilon) &= L(\phi_{cl} + \epsilon\tau\phi, \partial\phi_{cl} + \partial\epsilon\tau\phi + \epsilon\tau\partial\phi) \\ &\rightarrow \frac{\partial L(\phi_{cl}, \partial\phi_{cl})}{\partial\phi} \epsilon\tau + \frac{\partial L(\phi_{cl}, \partial\phi_{cl})}{\partial\partial\phi} [\partial\epsilon\tau\phi + \epsilon\tau\partial\phi]. \end{aligned} \quad (\text{A36})$$

The symmetry, Eq. (A35), indicates that ϵ is a cyclic coordinate and the equation of motion

$$\frac{\partial L(\epsilon, \partial\epsilon)}{\partial\epsilon} - \partial_\mu \frac{\partial L(\epsilon, \partial\epsilon)}{\partial\partial_\mu\epsilon} = 0. \quad (\text{A37})$$

shows that the current,

$$J^\mu = \frac{\partial L(\epsilon, \partial\epsilon)}{\partial\partial_\mu\epsilon} \quad (\text{A38})$$

defined up to a multiplicative constant as the generalized momentum of ϵ , is conserved. Notice that (i) we have an independent conserved current corresponding to each independent direction in the internal symmetry group and (ii) the conserved current is well defined up to a multiplicative constant only.

Examples:

1. Real scalar field: ϕ_a , $a = 1, \dots, n$, the symmetry group is $G = O(n)$,

$$\begin{aligned} L &= \frac{1}{2}(\partial\phi)^2 - V(\phi) \\ \delta\phi &= \epsilon^a \tau^a \phi, \quad \tau^a \in o(n) \\ J_\mu^a &= \partial_\mu \phi \tau^a \phi \end{aligned} \quad (\text{A39})$$

2. Complex scalar field: ϕ_a , $a = 1, \dots, n$, $G = \mathfrak{u}(n)$

$$\begin{aligned} L &= \partial\phi^\dagger \partial\phi - V(\phi) \\ \delta\phi &= i\epsilon^a \tau^a \phi, \quad \tau^a \in u(n) \\ J_\mu^a &= i\partial_\mu \phi^\dagger \tau^a \phi - i\phi^\dagger \tau^a \partial_\mu \phi \end{aligned} \quad (\text{A40})$$

3. Electromagnetic current: ϕ , $G = \mathfrak{so}(2) = SO(2)$, $\phi = \phi_1 + i\phi_2 = \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}$, $i = \begin{pmatrix} 0, -1 \\ 1, 0 \end{pmatrix}$

$$\begin{aligned} L &= \frac{1}{2}\partial_\mu \phi_1 \partial^\mu \phi_1 + \frac{1}{2}\partial_\mu \phi_2 \partial^\mu \phi_2 - m^2(\phi_1^2 + \phi_2^2) - V(\phi_1^2 + \phi_2^2) \\ J_\mu &= -(\partial_\mu \phi_1, \partial_\mu \phi_2) \begin{pmatrix} 0, -1 \\ 1, 0 \end{pmatrix} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} = -\partial_\mu \phi_2 \phi_1 + 2\partial_\mu \phi_1 \phi_2 = i(\partial_\mu \phi^\dagger \phi - \phi^\dagger \partial_\mu \phi) = -i\phi^\dagger \overleftrightarrow{\partial}_\mu \phi \end{aligned} \quad (\text{A41})$$

c. External Symmetries

The non-relativistic external symmetry group, the Galilean group consists of translations of the space-time and rotations of the space, a 4 + 3 dimensional continuous group. The relativistic




FIG. 10: Deformation of the space-time region.

Poincaré group is 10 dimensional, it is the direct sum of the translations and the Lorentz transformations in the space-time. We shall consider the conserved currents related to the translation invariance only for the sake of simplicity.

Let us rewrite the action in a space-time region V after an infinitesimal change of the coordinates, $x^\mu \rightarrow x'^\mu = x^\mu + \epsilon^\mu(x)$, $V \rightarrow V'$, cf. Fig. 10,

$$\begin{aligned} S[\phi] &= \int_V dx L(\phi(x), \partial\phi(x)) \\ &= \int_V \frac{dx}{\det(\delta_\nu^\mu + \partial_\nu \epsilon^\mu)} L(\phi(x - \epsilon), \partial\phi(x - \epsilon)) + \int_{V' - V} dx L(\phi(x), \partial\phi(x)), \end{aligned} \quad (\text{A42})$$

where the $\mathcal{O}(\epsilon^2)$ contributions are neglected in the second integral. The expansion in ϵ in the first integral, followed by a partial integration and the writing of the integral over $V' - V$ as a surface integral yields

$$\begin{aligned} S[\phi] &= \int_V dx \left[-\epsilon^\nu \partial_\mu \phi \left(\frac{\partial L(\phi, \partial\phi)}{\partial \phi} - \partial_\mu \frac{\partial L(\phi, \partial\phi)}{\partial \partial_\mu \phi} \right) - \partial_\nu \epsilon^\nu L(\phi, \partial\phi) \right] \\ &\quad + \int_{\partial V} dS_\mu \left[-\epsilon^\nu \partial_\mu \phi \frac{\partial L(\phi, \partial\phi)}{\partial \partial_\mu \phi} + \epsilon^\mu L(\phi, \partial\phi) \right], \end{aligned} \quad (\text{A43})$$

up to $\mathcal{O}(\epsilon)$ terms. The field configuration satisfies the equation of motion hence and consider a rigid shift of the space-time, $\epsilon(x) = \epsilon$ therefore the first line is vanishing. Since the left hand side is ϵ -independent for arbitrary space-time region, V , the energy-momentum tensor,

$$T^{\mu\nu} = \frac{\partial L}{\partial \partial_\nu \phi} \partial^\mu \phi - g^{\mu\nu} L \quad (\text{A44})$$

is conserved, $\partial_\mu T^{\mu\nu} = 0$ and the charge of the translation ϵ^ν ,

$$P^\mu = \int d^3x T^{0\mu}, \quad (\text{A45})$$

gives the energy-momentum vector. The energy-momentum tensor can be parametrized by

$$T^{\mu\nu} = \begin{pmatrix} \epsilon & c\mathbf{P} \\ \frac{1}{c}\mathbf{s} & \frac{1}{c}\sigma \end{pmatrix} \quad (\text{A46})$$

where

$$\begin{aligned}
\epsilon &= \text{energy density} \\
\mathbf{p} &= \text{momentum density} \\
\mathbf{S} &= \text{energy flux density} \\
\sigma^{jk} &= \text{momentum flux } p^k \text{ in the direction } j
\end{aligned} \tag{A47}$$

(c is restored).

The Lorentz symmetry leads to six conserved currents, 3 of which give the angular momentum and other three are the generators of the Lorentz boosts. The conservation of angular momentum can be used to prove that the energy-momentum tensor is symmetric, $T^{\mu\nu} = T^{\nu\mu}$ for bosonic field theories.

Appendix B: Parallel transport along a path

The expression of parallel transport, the solution to Eq. (37) is worked out in this Appendix. The solution can formally be written as

$$W_\gamma(y, x) = P \left[e^{-\int_x^y d\gamma^\mu A_\mu(\gamma)} \right] = P \left[e^{-\int_0^1 ds \frac{d\gamma^\mu(s)}{ds} A_\mu(\gamma(s))} \right], \tag{B1}$$

by means of the path ordered product of non-commuting objects defined along the path γ , defined as

$$P[A(s_A)B(s_B)] = \Theta(s_A - s_B)A(s_A)B(s_B) + \Theta(s_B - s_A)B(s_B)A(s_A). \tag{B2}$$

To see that we have the correct solution let us write first the integral in the exponent in Eq. (B1) as

$$\int_0^1 ds \frac{d\gamma^\mu(s)}{ds} A_\mu(\gamma(s)) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N \frac{d\gamma^\mu(s_j)}{ds} A_\mu(\gamma(s_j)) \tag{B3}$$

where $s_j = j/N$. The exponential function of an operator is defined by its Taylor-series,

$$\begin{aligned}
e^{-\frac{1}{N} \sum_{j=1}^N \frac{d\gamma^\mu(s_j)}{ds} A_\mu(\gamma(s_j))} &= 1 - \frac{1}{N} \sum_{j=1}^N \frac{d\gamma^\mu(s_j)}{ds} A_\mu(\gamma(s_j)) \\
&+ \frac{1}{N^2} \sum_{j_1, j_2=1}^N \frac{d\gamma^\mu(s_{j_1})}{ds} \frac{d\gamma^\mu(s_{j_2})}{ds} A_\mu(\gamma(s_{j_1})) A_\mu(\gamma(s_{j_2})) + \dots, \tag{B4}
\end{aligned}$$

and the path ordering applies term-by-term,

$$W_\gamma(y, x) = \lim_{N \rightarrow \infty} \left[1 - \frac{1}{N} \sum_{j=1}^N \frac{d\gamma^\mu(s_j)}{ds} A_\mu(\gamma(s_j)) + \frac{1}{N^2} \sum_{j_1, j_2=1}^N \frac{d\gamma^\mu(s_{j_1})}{ds} \frac{d\gamma^\mu(s_{j_2})}{ds} P[A_\mu(\gamma(s_{j_1})) A_\mu(\gamma(s_{j_2}))] + \dots \right]. \quad (\text{B5})$$

We would have

$$W_\gamma(y, x) = \lim_{N \rightarrow \infty} \prod_{j=1}^N e^{-\frac{1}{N} \frac{d\gamma^\mu(s_j)}{ds} A_\mu(\gamma(s_j))} \quad (\text{B6})$$

according to the well known rule $e^a e^b = e^{a+b}$, valid for numbers, without paying attention to the non-commutativity of the objects occurring in the product. But the path ordering places the contributions corresponding to higher j more to the left in the products and we find

$$W_\gamma(y, x) = \lim_{N \rightarrow \infty} e^{-\frac{1}{N} \frac{d\gamma^\mu(s_N)}{ds} A_\mu(\gamma(s_N))} \dots e^{-\frac{1}{N} \frac{d\gamma^\mu(s_1)}{ds} A_\mu(\gamma(s_1))} \quad (\text{B7})$$

by repeating the same resummation as in Eq. (B6). The path ordering succeeded in factorizing the dependence on the N -th division point to the integral at the very left of the product. The final step is the calculation of the partial derivative of W along the path,

$$\begin{aligned} \frac{1}{N} \frac{d\gamma^\mu}{ds} \partial_{y^\mu} W_\gamma(y, x) &= W_\gamma(y, x) - W_\gamma \left(y - \frac{1}{N} \frac{d\gamma^\mu(s_N)}{ds}, x \right) \\ &= \left[e^{-\frac{1}{N} \frac{d\gamma^\mu(s_N)}{ds} A_\mu(\gamma(s_N))} - \mathbb{1} \right] e^{-\frac{1}{N} \frac{d\gamma^\mu(s_{N-1})}{ds} A_\mu(\gamma(s_{N-1}))} \dots e^{-\frac{1}{N} \frac{d\gamma^\mu(s_1)}{ds} A_\mu(\gamma(s_1))} \\ &\approx -\frac{1}{N} \frac{d\gamma^\mu(s_N)}{ds} A_\mu(\gamma(s_N)) W_\gamma \left(y - \frac{1}{N} \frac{d\gamma^\mu(s_N)}{ds}, x \right) \\ &\approx -\frac{1}{N} \frac{d\gamma^\mu(1)}{ds} A_\mu(\gamma(1)) W_\gamma(y, x), \end{aligned} \quad (\text{B8})$$

which yields Eq. (37).

Appendix C: Higgs mechanism in scalar electrodynamics

First we discuss the spontaneous breakdown of the symmetry for a complex scalar field. Next the presence of massless modes are pointed out, finally the Higgs mechanism is presented in scalar electrodynamics.

1. Spontaneous symmetry breaking

Let us consider a scalar field theory, governed by the Lagrangian

$$L = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \mathcal{V}(\phi) \quad (\text{C1})$$

where

$$\mathfrak{V}(\phi) = \frac{m_0^2}{2}\phi^2 + \frac{g}{4}\phi^4, \quad (\text{C2})$$

which displays the symmetry $\phi \rightarrow -\phi$. The energy is given by

$$E = \int d^3x \left[\frac{1}{2}(\partial_0\phi)^2 + \frac{1}{2}(\nabla\phi)^2 + \mathfrak{V}(\phi) \right]. \quad (\text{C3})$$

A symmetry of the Lagrangian is not necessarily present in the solutions. The symmetry is called spontaneously broken if it is reflected by the Lagrangian but is violated by the "most important" field configuration, the one of having the lowest energy. For instance, the symmetry $\phi(x) \rightarrow -\phi(x)$ is broken spontaneously if the potential $\mathfrak{V}(\phi)$ has non-degenerate minima, for instance if $m_0^2 < 0$. In fact, if the initial conditions $\phi(t_i, \mathbf{x}) \sim \phi_0$, $\partial_0\phi(t_i, \mathbf{x}) \sim 0$ are used then the barrier of the potential energy does not allow the field to assume the symmetric, $\phi(x) = 0$, configuration.

To find the particle content of the they with spontaneously broken symmetry we write $\phi(x) = \phi_0 + \chi(x)$ where $\mathfrak{V}'(\phi) = 0$, $\mathfrak{V}''(\phi) > 0$, in particular

$$\phi_0 = \sqrt{\frac{-m_0^2}{g}}. \quad (\text{C4})$$

The Lagrangian for χ ,

$$L = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{1}{2}\mathfrak{V}''(\phi)\chi^2 + \mathcal{O}(\chi^3), \quad (\text{C5})$$

describes a field of mass $m^2 = -2m_0^2$ and the symmetry $\chi(x) \rightarrow -\chi(x)$ is not respected anymore.

2. Goldstone theorem

We now generalize the spontaneous breakdown of a discrete global symmetry to a continuous one by considering a complex scalar field whose dynamics is define by the Lagrangian

$$L = \partial_\mu\phi^*\partial^\mu\phi - \mathfrak{V}(\phi^*\phi) \quad (\text{C6})$$

with

$$\mathfrak{V}(\phi^*\phi) = m_0^2\phi^*\phi + \frac{g}{2}(\phi^*\phi)^2. \quad (\text{C7})$$

The Lagrangian remains invariant under the phase transformation $\phi(x) \rightarrow e^{i\alpha}\phi(x)$. The energy is now

$$E = \int d^3x [\partial_0\phi^*\partial_0\phi + \nabla\phi^*\nabla\phi + \mathfrak{V}(\phi^*\phi)]. \quad (\text{C8})$$

The potential with $m_0^2 > 0$ has a shape of a "Mexican hat", with degenerate minima at $\phi = e^{i\alpha}\phi_0$, where $\check{V}'(|\phi_0|^2) = 0$, $\check{V}''(|\phi_0|^2) > 0$. This suggest the parametrization

$$\phi(x) = [\phi_0 + \chi(x)]e^{i\alpha(x)} \quad (\text{C9})$$

of the complex field because the phase, parametrized by the field $\alpha(x)$ drops from the potential energy and has kinetic energy only. The Lagrangian for χ and α ,

$$L_m = \partial_\mu \chi \partial^\mu \chi + (\phi_0 + \chi)^2 \partial_\mu \alpha \partial^\mu \alpha - 2\check{V}''(\phi_0^2)\phi_0^2 \chi^2 + \mathcal{O}(\chi^3) \quad (\text{C10})$$

supports this idea, the field α has kinetic energy only and its normal modes have the dispersion relation, $\omega = \pm|\mathbf{p}|$, of a massless particle. For any spontaneously broken continuous symmetry the local, space-time dependent symmetry transformations define the Goldstone mode, having kinetic energy only and no potential one. As a result the Goldstone modes belong to massless particles.

The phase symmetry might be restored by the Goldstone modes since there is no potential energy barrier anymore. This in fact happens in any finite space volume. It is a non-trivial result that such a symmetry restoring modes slow down in the thermodynamical limit where the symmetry remains broken.

3. Higgs mechanism

Local gauge symmetry can not be broken spontaneously but the global component may break in the thermodynamical limit. To see what happens then we let the complex field interact with the electromagnetic field by minimal coupling, as given by the Lagrangian $L = L_{ED} + L_m$, where

$$L_{ED} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} \quad (\text{C11})$$

and

$$\begin{aligned} L_m &= (D_\mu \phi)^* D^\mu \phi + m^2 \phi^* \phi - \frac{g}{2}(\phi^* \phi)^2 \\ &= (\partial_\mu \phi^* + ieA_\mu \phi^*)(\partial^\mu \phi - ieA^\mu \phi) + m^2 \phi^* \phi - \frac{g}{2}(\phi^* \phi)^2 \\ &= \partial_\mu \phi^* \partial^\mu \phi + eA_\mu i(\phi^* \partial^\mu \phi - \partial^\mu \phi^* \phi) + e^2 A_\mu A^\mu \phi^* \phi + m^2 \phi^* \phi - \frac{g}{2}(\phi^* \phi)^2 \end{aligned} \quad (\text{C12})$$

We shall use the parametrization (C9) and choose the gauge $\alpha(x) = 0$ where

$$\begin{aligned} L_m &= \partial_\mu \chi \partial^\mu \chi + e^2 A_\mu A^\mu (\phi_0^2 + 2\phi_0 \chi + \chi^2) - \check{V}(\phi_0^2 + 2\phi_0 \chi + \chi^2) \\ &= \partial_\mu \chi \partial^\mu \chi + e^2 \phi_0^2 A_\mu A^\mu - 2g\phi_0^2 \chi^2 + \mathcal{O}(\chi^3) + \mathcal{O}(A^2 \chi) \end{aligned} \quad (\text{C13})$$

Therefore the Goldstone mode, an unphysical gauge mode can be eliminated by the help of gauge invariance and as a result the gauge field became massive,

$$M_A^2 = 2e^2\phi_0^2 = -\frac{2e^2}{g}m_0^2. \quad (\text{C14})$$

The surviving amplitude of the scalar field, the Higgs field has the mass

$$m_H^2 = 2g\phi_0^2 = -2m_0^2. \quad (\text{C15})$$

Appendix D: Scalar particle on a homogeneous non-Abelian magnetic field

We construct the particles, the normal modes of a free, scalar field ϕ with $S^{\mathfrak{su}(2)}$ local gauge invariance on a homogeneous magnetic gauge background field. The dynamics is governed by the Lagrangian

$$\begin{aligned} L_m &= (D_\mu\phi)^\dagger D^\mu\phi - m^2\phi^\dagger\phi \\ &= (\partial_\mu\phi^\dagger + g\phi^\dagger A_\mu)(\partial^\mu - gA^\mu)\phi - m^2\phi^\dagger\phi, \end{aligned} \quad (\text{D1})$$

where $A_\mu = A_\mu^a \frac{i}{2}\sigma^a$, in terms of the Pauli matrices σ^a and the background gauge field is chosen to be

$$A_\mu^a = H\delta^{a3}g_{\mu 3}x, \quad (\text{D2})$$

We seek the solution in factorized form,

$$\phi(t, x, y, z) = \psi(x)e^{-i\omega t + ik_y y + ik_z z}, \quad (\text{D6})$$

where the function $\psi(x)$ satisfies the equation

$$\left(-\partial_1^2 - gHxk_y\sigma^3 + \frac{1}{4}g^2H^2x^2\right)\psi = (\omega^2 - k_y^2 - k_z^2 - m^2)\psi \quad (\text{D7})$$

which can be written as a non-relativistic Schrödinger equation,

$$\left[-\frac{1}{2}\partial_1^2 + \frac{1}{2}\Omega^2\left(x - \frac{2k_y}{gH}\sigma^3\right)^2\right]\psi = \epsilon\psi, \quad (\text{D8})$$

with $\Omega = \frac{1}{2}gH$ and $\epsilon = \frac{\omega^2 - k_z^2 - m^2}{2}$. The solution $\psi = (\psi^+, \psi^-)$,

$$\psi_n^\pm(x) = H_n\left(x \mp \frac{2k_y}{gH}\right)e^{-\frac{1}{2}\Omega x^2} \quad (\text{D9})$$

describes a harmonic oscillator, centered at $x_0 = \pm \frac{2k_y}{gH}$, and the corresponding energy,

$$E_n(k_z) = \omega_n(k_z) = \pm\sqrt{(2n+1)\Omega + k_z^2 + m^2}, \quad (\text{D10})$$

depends on the discrete Landau-level quantum number n and the component of the wave vector in the direction of the magnetic field. The absence of k_y represents the continuous degeneracy of the Landau levels.

Appendix E: Gauge theory of the Poincaré group

The brief derivation of the Euler-Lagrange equations is presented here for the gauge theory formalism based on the Poincaré-group. The external space is the space-time as usual. The gauge group is chosen to be the symmetry of the local dynamics expe

Notice that this metric is homogeneous, space-time independent as required by the Equivalence Principle. Gauge transformations act as

$$\xi^a(x) \rightarrow \xi^a(x) + \Lambda^a_b(x)\xi^b(x) + \zeta^a(x) \quad (\text{E2})$$

where $\xi^a(x)$ is the internal space coordinate corresponding to the space-time point x , Λ^a_b denotes a Lorentz transformation matrix and ζ^a stands for translations. The gauge group, the Poincaré group, is the direct product of translations and Lorentz transformations, $P = T \times L$.

1. Covariant derivatives

We shall introduce the covariant derivative in the following steps. First we construct it for scalars $s(x)$, vectors $v^a(x)$ and tensors $t^{ab\dots}(x)$ of the internal Lorentz symmetry. Next we extend it for vectors or tensors defined by the space-time, $v^\mu(x)$, $t^{\mu\nu\dots}(x)$.

a. Vierbein

Let us start by recalling that the unique feature of gravity as gauge theory is that the internal space is the tangent space of the base manifold. Therefore there is a unique correspondence between infinitesimal translations in the internal and the external spaces and the dependence on x and ξ can be traded locally. A function $f(y)$ defined in the vicinity of x gives rise the function

$$f_x(\xi) = f(x + \Delta x) \quad (\text{E3})$$

defined in the vicinity of the origin of the Lorentz space where

$$\xi^a = e^a_\mu \Delta x^\mu. \quad (\text{E4})$$

The matrix e^a_μ relating the directions in the two spaces,

$$e^a_\mu = \frac{\partial \xi^a}{\partial x^\mu}, \quad (\text{E5})$$

is called vierbein. The relation

$$e^a_\mu \partial_a s = \frac{\partial s}{\partial x^\mu} = \partial_\mu s \quad (\text{E6})$$

follows in an obvious manner and allows to represent the functions over the space-time locally as functions over the tangent space, according to the intuitive role played by the coordinates in the Equivalence Principle.

The inverse of the transformation (E5) is

$$e_a^\mu = \frac{\partial x^\mu}{\partial \xi^a}, \quad (\text{E7})$$

and the metric tensor is given by

$$g^{\mu\nu}(x) = e_a^\mu(x)\eta^{ab}e_b^\nu(x). \quad (\text{E8})$$

Thus we have the identities

$$e_a^\mu e_\nu^a = \delta_\nu^\mu, \quad e_a^\mu e_\mu^b = \delta_a^b, \quad (\text{E9})$$

and any vector or tensor can be represented as a Lorentz or world vector or tensor as in Eq. (E6), eg.

$$v^\mu = \frac{\partial x^\mu}{\partial \xi^a} v^a = e_a^\mu v^a, \quad v^a = \frac{\partial \xi^a}{\partial x^\mu} v^\mu = e_\mu^a v^\mu. \quad (\text{E10})$$

Note that the invariant integral measure can be written as

$$\sqrt{-\det g_{\mu\nu}} d^4x = \det(e_\mu^a) d^4x. \quad (\text{E11})$$

b. Holonomic and anholonomic vector fields

The Equivalence Principle assures that the gravitation interaction can locally be eliminated by a suitable choice of the coordinate system which is realised by the internal space. This special coordinate system can be extended in a finite regions in the absence of gravitational field only. What is the condition on four world vector fields $e_a^\mu(x)$ with $a = 0, 1, 2, 3$ for the existence of a coordinate systems with these coordinate axes?

The solution of the differential equation

$$\frac{\partial x^\mu(\xi)}{\partial \xi} = e^\mu(x) \quad (\text{E12})$$

can be written as

$$x^\mu(\xi) = e^{\xi \frac{\partial}{\partial \xi}} x^\mu(0). \quad (\text{E13})$$

We are now looking for the coordinates, defined by the coordinate axes

$$\frac{\partial x^\mu(\xi)}{\partial \xi^a} = e_a^\mu(x). \quad (\text{E14})$$

Let us suppose that the world vectors $e_a^\mu(x)$ are independent and the derivatives $\partial_a = \frac{\partial}{\partial \xi^a}$ commute,

$$\begin{aligned} [\partial_a, \partial_b] &= [e_a^\mu \partial_\mu, e_b^\nu \partial_\nu] \\ &= (e_a^\mu \partial_\mu e_b^\nu - e_b^\mu \partial_\mu e_a^\nu) \partial_\nu = 0. \end{aligned} \quad (\text{E15})$$

Then the coordinates can locally be introduced for example by

$$x^\mu(\xi) = e^{\xi^0 \frac{\partial}{\partial \xi^0}} e^{\xi^1 \frac{\partial}{\partial \xi^1}} e^{\xi^2 \frac{\partial}{\partial \xi^2}} e^{\xi^3 \frac{\partial}{\partial \xi^3}} x^\mu(\xi)_{\xi=0}. \quad (\text{E16})$$

In fact, by varying ξ around zero we cover all four dimensions and

$$\begin{aligned} \partial_a x^\mu(0) &= \partial_a e^{\xi^0 \frac{\partial}{\partial \xi^0}} e^{\xi^1 \frac{\partial}{\partial \xi^1}} e^{\xi^2 \frac{\partial}{\partial \xi^2}} e^{\xi^3 \frac{\partial}{\partial \xi^3}} x^\mu(\xi)_{\xi=0} \\ &= \partial_a \left(1 + \xi^a \frac{\partial}{\partial \xi^a} \right) x^\mu(\xi)_{\xi=0} \\ &= e_a^\mu(x(0)). \end{aligned} \quad (\text{E17})$$

When the vectors e_a^μ are independent only but the commutators are non-vanishing then ξ^a is already an admissible coordinate system but the coordinate axes are different than the vectors e_a^μ .

Conversely, let us suppose that the the solution of the system of differential equations lead to admissible coordinates. The invertibility of the coordinate transformation requires the independence of the world vectors e_a^μ and the commutativity of the derivatives, ie. the symmetry of partial derivatives of continuously derivable functions is trivial in the coordinate system ξ^a .

The vectors e_a^μ are called holonomic if they are the directional vectors of a local coordinate system. Therefore the vierbein defines anholonomic vectors in the presence of gravitational field.

c. Local Lorentz transformations

The covariant derivative involves a generator valued vector field. The gauge group is a direct product, $P = T \times L$, therefore there will be separate gauge fields for translations and Lorentz transformations. Due to continuity requirements the proper Lorentz group is generated by these generators only, discrete inversions will be left out.

Local Lorentz transformations generate space-dependent orientation for the orthogonal Lorentz coordinate basis and lead to the covariant derivative

$$D_\mu^{(L)} = \frac{\partial}{\partial x^\mu} + \omega_\mu \quad (\text{E18})$$

where the affine connection, $\omega_{ab\mu} = \eta_{ac}(\omega_\mu)^c_b = \eta_{ac}\omega^c_{b\mu}$, $\omega_{ab\mu} = -\omega_{ba\mu}$ is a generator of the special Lorentz group in the fundamental representation (E2). The connection acts in the internal space

only, ie. on Lorentz tensors $T_{a,b,\dots}$, eg.

$$D_\mu^{(L)}v^b = \partial_\mu v^b + \omega^b_{c\mu}v^c \quad (\text{E19})$$

and considers the world tensors $T_{\mu\nu\dots}$ as scalars,

$$D_\mu^{(L)}v^\mu = \partial_\mu v^\mu. \quad (\text{E20})$$

In order to preserve the Lorentzian scalar product by parallel transport,

$$D_\mu^{(L)}(v_b u^b) = \partial_\mu(v_b u^b) \quad (\text{E21})$$

we define

$$D_\mu^{(L)}v_b = \partial_\mu v_b - v_c \omega^c_{b\mu}. \quad (\text{E22})$$

Notice the vanishing of the covariant derivative of the internal space metric,

$$\begin{aligned} D_\mu^{(i)}\eta^{ab} &= \omega^a_{c\mu}\eta^{cb} + \omega^b_{c\mu}\eta^{ac} \\ &= \omega^{ab}_\mu + \omega^{ba}_\mu \\ &= 0. \end{aligned} \quad (\text{E23})$$

d. Local translations

Local translations generate space-time dependent shift of the Lorentz coordinate system which in turn induces a shift in the space-time due to the fact that the internal space is the tangent space of the space-time. The corresponding covariant derivative is

$$D_\mu^{(T)}s = \delta_\mu^a \partial_a s + t_\mu^a \partial_a s = e_\mu^a \partial_a s \quad (\text{E24})$$

for a scalar function $s(x)$ with $t_\mu^a(x)$ as gauge field. The first term on the right hand side implements the infinitesimal shift $\Delta x^\mu D_\mu^{(T)}$ in the internal space which corresponds to an infinitesimal shift $\Delta x^\mu \partial_\mu$ in the external space and the second term compensates for the difference of the position of the origin of the internal coordinate system at the space-time points x and $x + \Delta x$. Thus the vierbein, satisfying Eq. (E6) is

$$e_\mu^a = \delta_\mu^a + t_\mu^a. \quad (\text{E25})$$

e. Full local Poincare group

The full internal symmetry is incorporated into the covariant derivative $D_\mu^{(i)}$ acts as $D_\mu^{(L)}$,

$$\begin{aligned} D_\mu^{(i)} s &= D_\mu^{(T)} s = D_\mu^{(L)} s = \partial_\mu s \\ D_\mu^{(i)} v^b &= \delta_\mu^a \partial_a v^b + t_\mu^a \partial_a v^b + \omega_{c\mu}^b v_c = \partial_\mu v^b + \omega_{c\mu}^b v_c = D_\mu^{(L)} v^b \\ D_\mu^{(i)} v_b &= \delta_\mu^a \partial_a v_b + t_\mu^a \partial_a v_b - v_c \omega_{b\mu}^c = \partial_\mu v_b - v_c \omega_{b\mu}^c = D_\mu^{(L)} v_b \end{aligned} \quad (\text{E26})$$

on scalars and Lorentz tensors and treating world vectors and tensors as scalars because ∂_μ can be considered as the covariant derivative for local translations.

It is an unusual feature that one can introduce derivatives in the internal space directions, ∂_a . The corresponding covariant derivative is

$$D_a^{(i)} = e_a^\mu D_\mu^{(i)} = e_a^\mu (\partial_\mu + \omega_\mu). \quad (\text{E27})$$

Notice the natural relation

$$D_a^{(i)} v = \partial_a v + e_a^\mu t_\mu^b \partial_b v = \partial_a v + e_a^\mu (e_\mu^b - \delta_\mu^b) \partial_b v = 2\partial_a v - e_a^\mu \delta_\mu^b \partial_b v = \partial_a v. \quad (\text{E28})$$

f. World vectors and tensors

We could stop at this stage and start to work out the gauge theory for the Poincaré group by means of the covariant derivative $D^{(i)}$. The drawback would be to use the internal Minkowski space vectors v^a or tensors $t^{ab\dots}$ only. This is obviously an artificial constraint because the coordinates x^μ lead naturally to world vectors v^μ or tensors $t^{\mu\nu\dots}$. In order to be construct covariant equations for these vectors or tensors we need the covariant derivative for the $GL(4)$ gauge theory, controlling the effects of coordinate transformations induced by the application of the local Poincaré group in the internal space. The usual affine connection Γ_μ , introduced in differential geometry, realizes this covariant derivative

$$D_\mu^{(e)} = \partial_\mu + \Gamma_\mu \quad (\text{E29})$$

by acting on world scalars, vectors and tensors, eg.

$$\begin{aligned} D_\mu^{(e)} s &= \partial_\mu s \\ D_\mu^{(e)} v^\nu &= \partial_\mu v^\nu + \Gamma_{\rho\mu}^\nu v^\rho \\ D_\mu^{(e)} v_\nu &= \partial_\mu v_\nu - v_\rho \Gamma_{\rho\mu}^\nu. \end{aligned} \quad (\text{E30})$$

The field strength tensor corresponding to internal directions is

$$\begin{aligned}
F_{ab} &= [D_a, D_b] = [e_a^\mu D_\mu, e_b^\nu D_\nu] \\
&= e_a^\mu e_b^\nu [D_\mu, D_\nu] + e_a^\mu [D_\mu, e_b^\nu] D_\nu + e_b^\nu [e_a^\mu, D_\nu] D_\mu + [e_a^\mu, e_b^\nu] D_\nu D_\mu \\
&= e_a^\mu e_b^\nu F_{\mu\nu} + e_a^\mu [D_\mu, e_b^\nu] D_\nu + e_b^\nu [e_a^\mu, D_\nu] D_\mu \\
&= e_a^\mu e_b^\nu F_{\mu\nu} + e_a^\mu D_\mu e_b^\nu D_\nu - e_a^\mu e_b^\nu D_\mu D_\nu + e_b^\nu e_a^\mu D_\nu D_\mu - e_b^\nu D_\nu e_a^\mu D_\mu \\
&= e_a^\mu e_b^\nu F_{\mu\nu} + e_a^\mu (D_\mu e_b^\nu) D_\nu - e_b^\nu (D_\nu e_a^\mu) D_\mu \\
&= e_a^\mu e_b^\nu F_{\mu\nu} + S_{ab}^\mu D_\mu
\end{aligned} \tag{E39}$$

with

$$\begin{aligned}
S_{ab}^\mu &= e_a^\nu D_\nu e_b^\mu - e_b^\nu D_\nu e_a^\mu \\
&= D_a e_b^\mu - D_b e_a^\mu \\
&= e_a^\nu (\partial_\nu e_b^\mu - e_c^\mu \omega_{b\nu}^c + \Gamma_{\rho\nu}^\mu e_b^\rho) - e_b^\nu (\partial_\nu e_a^\mu - e_c^\mu \omega_{a\nu}^c + \Gamma_{\rho\nu}^\mu e_a^\rho)
\end{aligned} \tag{E40}$$

being the torsion tensor. The translational part of D_a defines the field strength tensor

$$t_{ab}^\mu = \partial_a e_b^\mu - \partial_b e_a^\mu, \tag{E41}$$

cf. Eq.(E15), because

$$\begin{aligned}
[\partial_a, \partial_b] &= [e_a^\mu \partial_\mu, e_b^\nu \partial_\nu] \\
&= e_a^\mu [\partial_\mu, e_b^\nu] \partial_\nu + e_b^\nu [e_a^\mu, \partial_\nu] \partial_\mu + [e_a^\mu, e_b^\nu] \partial_\nu \partial_\mu \\
&= e_a^\mu [\partial_\mu, e_b^\nu] \partial_\nu + e_b^\nu [e_a^\mu, \partial_\nu] \partial_\mu \\
&= e_a^\mu \partial_\mu e_b^\nu \partial_\nu - e_a^\mu e_b^\nu \partial_\mu \partial_\nu + e_b^\nu e_a^\mu \partial_\nu \partial_\mu - e_b^\nu \partial_\nu e_a^\mu \partial_\mu \\
&= e_a^\mu (\partial_\mu e_b^\nu) \partial_\nu - e_b^\nu (\partial_\nu e_a^\mu) \partial_\mu \\
&= t_{ab}^\mu \partial_\mu.
\end{aligned} \tag{E42}$$

3. Variational equations

Let us suppose that the action can be written as

$$S = S_g[e, \omega] + S_m[\phi, e, \omega] \tag{E43}$$

where $S_m[\phi, e, \omega]$ controls the dynamics of the matter field, denoted generically by ϕ on a given geometry, specified by the fields e and ω . For a scalar field we may have

$$S_m[\phi, e, \omega] = \int dx E \left[\frac{1}{2} D_\mu^{(e)} \phi D^{(e)\mu} \phi - V(\phi) \right] \tag{E44}$$

where

$$E = \det e_{\mu}^a \quad (\text{E45})$$

and for a fermion

$$S_m[\bar{\psi}, \psi, e, \omega] = \int dx E \bar{\psi} i \gamma^a \underbrace{e_a^{\mu} (\partial_{\mu} + \omega_{\mu}^{ab} \tau_{ab})}_{D_a} \psi \quad (\text{E46})$$

with $\tau_{ab} = -\frac{1}{4}[\gamma_a, \gamma_b]$. The gravitational action is chosen to be of the Einstein-Hilbert type,

$$S_g[e, \omega] = -\frac{1}{16\pi G} \int dx ER. \quad (\text{E47})$$

a. Variation of ω

The scalar curvature can be written as

$$\begin{aligned} R &= \eta^{bc} F_{bac}^a \\ &= \eta^{bd} e_a^{\mu} e_d^{\nu} (\partial_{\mu} \omega_{b\nu}^a - \partial_{\nu} \omega_{b\mu}^a + \omega_{c\mu}^a \omega_{b\nu}^c - \omega_{c\nu}^a \omega_{b\mu}^c) \\ &= \eta^{bd} (e_a^{\mu} e_d^{\nu} - e_a^{\nu} e_d^{\mu}) (\partial_{\mu} \omega_{b\nu}^a + \omega_{c\mu}^a \omega_{b\nu}^c) \\ &= \eta^{bd} T_{ad}^{\mu\nu} (\partial_{\mu} \omega_{b\nu}^a + \omega_{c\mu}^a \omega_{b\nu}^c) \end{aligned} \quad (\text{E48})$$

where

$$T_{ab}^{\mu\nu} = e_a^{\mu} e_b^{\nu} - e_a^{\nu} e_b^{\mu}. \quad (\text{E49})$$

The variation of the Lorentz connection gives the equation

$$\begin{aligned} 0 &= -16\pi G \frac{\delta S}{\delta \omega_{b\mu}^a} = E \eta^{cd} T_{ad}^{\mu\nu} \omega_{c\nu}^b + E \eta^{bd} T_{cd}^{\nu\mu} \omega_{a\nu}^c - \partial_{\nu} (E \eta^{bd} T_{ad}^{\nu\mu}) \\ &= E T_a^{\mu\nu c} \omega_{c\nu}^b + E T_c^{\nu\mu b} \omega_{a\nu}^c - \partial_{\nu} (E T_a^{\nu\mu b}) \\ &= -E \omega_{c\nu}^b T_a^{\nu\mu c} + E T_c^{\nu\mu b} \omega_{a\nu}^c - \partial_{\nu} (E T_a^{\nu\mu b}) \\ &= -D_{\nu}^{(i)} (E T_a^{\nu\mu b}) \end{aligned} \quad (\text{E50})$$

The relation

$$\partial_{\mu} E = \partial_{\mu} e^{\text{tr} \ln e} = e^{\text{tr} \ln e} \text{tr} e^{-1} \partial_{\mu} e = E e_a^{\nu} \partial_{\nu} e_{\mu}^a \quad (\text{E51})$$

allows us to write it in the form

$$\begin{aligned} 0 &= E T_a^{\nu\mu b} e_c^{\rho} \partial_{\rho} e_{\nu}^c + E \partial_{\nu} T_a^{\nu\mu b} + E \omega_{c\nu}^b T_a^{\nu\mu c} - E T_c^{\nu\mu b} \omega_{a\nu}^c \\ &= E [T_a^{\nu\mu b} e_c^{\rho} \partial_{\rho} e_{\nu}^c + \partial_{\nu} T_a^{\nu\mu b} + \omega_{c\nu}^b T_a^{\nu\mu c} - T_c^{\nu\mu b} \omega_{a\nu}^c] \\ &= E [T_a^{\nu\mu b} e_c^{\rho} \partial_{\rho} e_{\nu}^c + D_{\nu}^{(i)} T_a^{\nu\mu b}] \end{aligned} \quad (\text{E52})$$

or

$$-T_a^{\nu\mu b} e_c^\rho \partial_\rho e_\nu^c = \partial_\nu T_a^{\nu\mu b} + \omega_{c\nu}^b T_a^{\nu\mu c} - T_c^{\nu\mu b} \omega_{a\nu}^c \quad (\text{E53})$$

which in turn gives

$$\begin{aligned} T_{ab}^{\nu\mu} \partial_\rho e_c^\rho e_\nu^c &= -T_{ab}^{\nu\mu} e_c^\rho \partial_\rho e_\nu^c \\ &= \partial_\nu T_{ab}^{\nu\mu} + \omega_{bc\nu} T_a^{\nu\mu c} - T_{cb}^{\nu\mu} \omega_{a\nu}^c \\ &= \partial_\nu T_{ab}^{\nu\mu} - \omega_{cb\nu} T_a^{\nu\mu c} - T_{cb}^{\nu\mu} \omega_{a\nu}^c \\ &= \partial_\nu T_{ab}^{\nu\mu} - \omega_{b\nu}^c T_{ac}^{\nu\mu} - T_{cb}^{\nu\mu} \omega_{a\nu}^c \\ &= D_\nu T_{ab}^{\nu\mu}. \end{aligned} \quad (\text{E54})$$

In order to solve the equation

$$T_{ab}^{\nu\mu} \partial_\rho e_c^\rho e_\nu^c = \partial_\nu T_{ab}^{\nu\mu} - \omega_{b\nu}^c T_{ac}^{\nu\mu} - T_{cb}^{\nu\mu} \omega_{a\nu}^c \quad (\text{E55})$$

for the Lorentz connection we write

$$\begin{aligned} (e_a^\nu e_b^\mu - e_a^\mu e_b^\nu) \partial_\rho e_c^\rho e_\nu^c &= \partial_\nu (e_a^\nu e_b^\mu - e_a^\mu e_b^\nu) - \omega_{b\nu}^c (e_a^\nu e_c^\mu - e_a^\mu e_c^\nu) - (e_c^\nu e_b^\mu - e_c^\mu e_b^\nu) \omega_{a\nu}^c \\ e_b^\mu \partial_\rho e_a^\rho - e_a^\mu \partial_\rho e_b^\rho &= \partial_\nu (e_a^\nu e_b^\mu - e_a^\mu e_b^\nu) + \omega_{bc}^c e_a^\mu - \omega_{ac}^c e_b^\mu + 2\omega_{ab}^\mu \end{aligned} \quad (\text{E56})$$

and we find

$$2\omega_{ab}^\mu + \omega_{bc}^c e_a^\mu - \omega_{ac}^c e_b^\mu = t_{ab}^\mu \quad (\text{E57})$$

where

$$t_{ab}^\mu = \partial_\nu (e_a^\nu e_b^\mu - e_a^\mu e_b^\nu) - e_b^\mu \partial_\rho e_a^\rho + e_a^\mu \partial_\rho e_b^\rho \quad (\text{E58})$$

It will be more useful to have Lorentz indices only for the connection, therefore we multiply Eq. (E57) by e_μ^d

$$2\omega_{ab}^d + \omega_{bc}^c \delta_a^d - \omega_{ac}^c \delta_b^d = t_{ab}^d. \quad (\text{E59})$$

To find the second and third terms on the left hand side we contract the indices a and d and find

$$\omega_{bc}^c = \frac{1}{3} t_{cb}^c \quad (\text{E60})$$

by recalling the antisymmetry of the spacial Lorentz group generators, $\omega_{abc} = -\omega_{bac}$. This result leads us to the solution

$$\omega_{ab}^d = \frac{1}{6} t_{ca}^c \delta_b^d - \frac{1}{6} t_{cb}^c \delta_a^d + \frac{1}{2} t_{ab}^d \quad (\text{E61})$$

where

$$\begin{aligned}
t^c_{ab} &= e^c_\mu \partial_\nu (e^\nu_a e^\mu_b - e^\mu_a e^\nu_b) - \delta^c_b \partial_\rho e^\rho_a + \delta^c_a \partial_\rho e^\rho_b \\
&= e^c_\mu \partial_\nu e^\nu_a e^\mu_b - e^c_\mu \partial_\nu e^\mu_a e^\nu_b + e^c_\mu e^\nu_a \partial_\nu e^\mu_b - e^c_\mu e^\mu_a \partial_\nu e^\nu_b - \delta^c_b \partial_\rho e^\rho_a + \delta^c_a \partial_\rho e^\rho_b \\
&= \delta^c_b \partial_\nu e^\nu_a - e^c_\mu e^\nu_b \partial_\nu e^\mu_a + e^c_\mu e^\nu_a \partial_\nu e^\mu_b - \delta^c_a \partial_\nu e^\nu_b - \delta^c_b \partial_\rho e^\rho_a + \delta^c_a \partial_\rho e^\rho_b \\
&= e^c_\mu e^\nu_a \partial_\nu e^\mu_b - e^c_\mu e^\nu_b \partial_\nu e^\mu_a \\
&= e^c_\mu (\partial_a e^\mu_b - \partial_b e^\mu_a) \\
&= e^c_\mu t^\mu_{ab}.
\end{aligned} \tag{E62}$$

is given in terms of the translational field strength tensor introduced in Eq. (E41). Its contracted expression,

$$\begin{aligned}
t^b_{ba} &= e^b_\mu (\partial_b e^\mu_a - \partial_a e^\mu_b) \\
&= \partial_\mu e^\mu_a - e^b_\mu e^\nu_a \partial_\nu e^\mu_b,
\end{aligned} \tag{E63}$$

inserted in Eq. (E61) gives

$$\begin{aligned}
\omega^c_{ab} &= \frac{1}{6} t^d_{da} \delta^c_b - \frac{1}{6} t^d_{db} \delta^c_a + \frac{1}{2} t^c_{ab} \\
&= \frac{1}{6} (\partial_\mu e^\mu_a - e^d_\mu e^\nu_a \partial_\nu e^\mu_d) \delta^c_b - \frac{1}{6} (\partial_\mu e^\mu_b - e^d_\mu e^\nu_b \partial_\nu e^\mu_d) \delta^c_a + \frac{1}{2} e^c_\mu e^\nu_a \partial_\nu e^\mu_b - \frac{1}{2} e^c_\mu e^\nu_b \partial_\nu e^\mu_a
\end{aligned} \tag{E64}$$

and

$$\begin{aligned}
\omega^c_{a\mu} &= \frac{1}{6} e^b_\mu \partial_\rho e^\rho_a \delta^c_b - \frac{1}{6} e^b_\mu e^d_\rho e^\nu_a \partial_\nu e^\rho_d \delta^c_b - \frac{1}{6} e^b_\mu \partial_\rho e^\rho_b \delta^c_a + \frac{1}{6} e^b_\mu e^d_\rho e^\nu_b \partial_\nu e^\rho_d \delta^c_a + \frac{1}{2} e^b_\mu e^c_\rho e^\nu_a \partial_\nu e^\rho_b - \frac{1}{2} e^b_\mu e^c_\rho e^\nu_b \partial_\nu e^\rho_a \\
&= \frac{1}{6} e^c_\mu \partial_\rho e^\rho_a - \frac{1}{6} e^c_\mu e^d_\rho e^\nu_a \partial_\nu e^\rho_d - \frac{1}{6} \delta^c_a e^b_\mu \partial_\rho e^\rho_b + \frac{1}{6} \delta^c_a e^d_\rho \partial_\mu e^\rho_d + \frac{1}{2} e^b_\mu e^c_\rho e^\nu_a \partial_\nu e^\rho_b - \frac{1}{2} e^c_\rho \partial_\mu e^\rho_a \\
&= \frac{1}{6} e^c_\mu e^b_\rho \partial_b e^\rho_a - \frac{1}{6} e^c_\mu e^d_\rho \partial_a e^\rho_d - \frac{1}{6} \delta^c_a e^b_\mu e^d_\rho \partial_d e^\rho_b + \frac{1}{6} \delta^c_a e^d_\rho e^b_\mu \partial_b e^\rho_d + \frac{1}{2} e^b_\mu e^c_\rho \partial_a e^\rho_b - \frac{1}{2} e^c_\rho e^b_\mu \partial_b e^\rho_a \\
&= \frac{1}{6} e^c_\mu e^b_\rho t^\rho_{ba} + \frac{1}{6} \delta^c_a e^b_\mu e^d_\rho t^\rho_{bd} + \frac{1}{2} e^b_\mu e^c_\rho t^\rho_{ab} \\
&= \frac{1}{6} (e^c_\mu \delta^d_a e^b_\rho + \delta^c_a e^b_\mu e^d_\rho - 3e^c_\rho \delta^d_a e^b_\mu) t^\rho_{bd}.
\end{aligned} \tag{E65}$$

Notice that the affine connection of the external space covariant derivative does not appear in this equation because the scalar curvature (E48) is expressed in terms of the field strength tensor of ω_μ .

b. Variation of e

We consider vierbein components e^a_μ as independent variables and the variation of

$$e^a_\mu e^\mu_a = \delta^a_a \tag{E66}$$

gives

$$\delta e_a^\nu e_\mu^a + e_a^\nu \delta e_\mu^a = 0 \quad (\text{E67})$$

and

$$\delta e_b^\nu = -e_b^\mu e_a^\nu \delta e_\mu^a. \quad (\text{E68})$$

The variational equation for the vierbein which appears in the tensor $T_{ab}^{\mu\nu}$, given by Eq. (E49) and in the determinant E is

$$0 = -16\pi G \frac{\delta S}{\delta e_\rho^b} = 16\pi G \frac{\delta S}{\delta e_c^\kappa} e_b^\kappa e_c^\rho = 16\pi G \left(E \frac{\delta R}{\delta T_{ad}^{\mu\nu}} \frac{\delta T_{ad}^{\mu\nu}}{\delta e_c^\kappa} + \frac{\delta E}{\delta e_c^\kappa} R \right) e_b^\kappa e_c^\rho \quad (\text{E69})$$

what we can write by means of Eqs. (E48) and

$$\delta E = \delta e^{\text{tr} \ln e} = e^{\text{tr} \ln e} \text{tr} e^{-1} \delta e = E e_a^\mu \delta e_\mu^a = -E e_\mu^a \delta e_a^\mu \quad (\text{E70})$$

as

$$\begin{aligned} 0 &= \eta^{bd} (\partial_\mu \omega_{b\nu}^a + \omega_{e\mu}^a \omega_{b\nu}^e) (\delta_\kappa^\mu \delta_a^c e_d^\nu + e_a^\mu \delta_\kappa^\nu \delta_d^c - \delta_\kappa^\nu \delta_a^c e_d^\mu - e_a^\nu \delta_\kappa^\mu \delta_d^c) - e_\kappa^c R \\ &= (\partial_\mu \omega_{\nu}^{ad} + \omega_{e\mu}^a \omega_{\nu}^{ed}) (\delta_\kappa^\mu \delta_a^c e_d^\nu + e_a^\mu \delta_\kappa^\nu \delta_d^c - \delta_\kappa^\nu \delta_a^c e_d^\mu - e_a^\nu \delta_\kappa^\mu \delta_d^c) - e_\kappa^c R \\ &= (\partial_\kappa \omega_{\nu}^{cd} + \omega_{e\kappa}^c \omega_{\nu}^{ed}) e_d^\nu + (\partial_\mu \omega_{\kappa}^{ac} + \omega_{e\mu}^a \omega_{\kappa}^{ec}) e_a^\mu - (\partial_\mu \omega_{\kappa}^{cd} + \omega_{e\mu}^c \omega_{\kappa}^{ed}) e_d^\mu \\ &\quad - (\partial_\kappa \omega_{\nu}^{ac} + \omega_{e\kappa}^a \omega_{\nu}^{ec}) e_a^\nu - e_\kappa^c R \end{aligned} \quad (\text{E71})$$

We contract the last equation with e_b^κ and write

$$\begin{aligned} 0 &= (\partial_\kappa \omega_{\nu}^{cd} + \omega_{e\kappa}^c \omega_{\nu}^{ed}) e_d^\nu e_b^\kappa + (\partial_\mu \omega_{\kappa}^{ac} + \omega_{e\mu}^a \omega_{\kappa}^{ec}) e_a^\mu e_b^\kappa \\ &\quad - (\partial_\mu \omega_{\kappa}^{cd} + \omega_{e\mu}^c \omega_{\kappa}^{ed}) e_d^\mu e_b^\kappa - (\partial_\kappa \omega_{\nu}^{ac} + \omega_{e\kappa}^a \omega_{\nu}^{ec}) e_a^\nu e_b^\kappa - \delta_b^c R \\ &= 2(e_d^\nu e_b^\kappa - e_d^\kappa e_b^\nu) (\partial_\kappa \omega_{\nu}^{cd} + \omega_{e\kappa}^c \omega_{\nu}^{ed}) - \delta_b^c R. \end{aligned} \quad (\text{E72})$$

The Ricci tensor

$$\begin{aligned} R_{bc} &= e_c^\nu e_a^\mu F_{b\mu\nu}^a \\ &= e_c^\nu e_a^\mu (\partial_\mu \omega_{b\nu}^a - \partial_\nu \omega_{b\mu}^a + \omega_{c\mu}^a \omega_{b\nu}^c - \omega_{c\nu}^a \omega_{b\mu}^c) \\ &= (e_c^\nu e_a^\mu - e_c^\mu e_a^\nu) (\partial_\mu \omega_{b\nu}^a + \omega_{c\mu}^a \omega_{b\nu}^c) \\ &= T_{ca}^{\nu\mu} (\partial_\mu \omega_{b\nu}^a + \omega_{c\mu}^a \omega_{b\nu}^c) \end{aligned} \quad (\text{E73})$$

gives finally the Einstein equation

$$R_{ab} - \frac{1}{2} \eta_{ab} R = 0 \quad (\text{E74})$$

c. *Affine connection for world vectors*

The affine connection Γ_μ appearing in the covariant derivative $D^{(e)}$ has not been included in the action (E47) and it will be determined by a non-dynamical principle. The parallel transport of a vector v^μ during a displacement δx^μ , expressed by the equation

$$\delta x^\nu D_\nu^{(e)} v^\mu = 0 \quad (\text{E75})$$

must be equivalent with the similar equation expressing the parallel transport of the vector v^a ,

$$\delta x^\nu D_\nu^{(i)} v^a = 0. \quad (\text{E76})$$

The covariant condition for the equivalence of the two parallel transports, valid for arbitrary vector field v^a , is

$$D_\nu^{(e)} e_a^\mu v^a = e_a^\mu D_\nu^{(i)} v^a. \quad (\text{E77})$$

The combination $e_a^\mu v^a$ is an internal space scalar,

$$D_\nu^{(e)} e_a^\mu v^a = D_\nu e_a^\mu v^a \quad (\text{E78})$$

thus we have

$$D_\nu e_a^\mu v^a = e_a^\mu D_\nu v^a, \quad (\text{E79})$$

or

$$D_\nu e_a^\mu = 0 \quad (\text{E80})$$

which leads to metric admissibility,

$$D_\nu e_a^\mu \eta^{ab} e_b^\kappa = D_\nu g^{\mu\kappa} = 0 \quad (\text{E81})$$

The affine connection, Γ , can easily be obtained by solving Eq. (E80),

$$D_\nu e_a^\mu = D_\nu^{(i)} e_a^\mu + \Gamma_{\rho\nu}^\mu e_a^\rho = 0 \quad (\text{E82})$$

with the result

$$\begin{aligned} \Gamma_{\rho\nu}^\mu &= -e_\rho^a D_\nu e_a^\mu \\ &= e_a^\mu D_\nu e_\rho^a \\ &= e_a^\mu \partial_\nu e_\rho^a + e_a^\mu \omega_{\nu b}^a e_\rho^b \neq \Gamma_{\nu\rho}^\mu. \end{aligned} \quad (\text{E83})$$