


^{89}Zr-labeled radioimmunoconjugates with bispidine ligands	
	Equipe de Synthèse pour l'Analyse (SynPA), Université de Strasbourg, CNRS, IPHC UMR 7178, 67000 Strasbourg Directrice de thèse : Aline Nonat (aline.nonat@unistra.fr)
	

Keywords: coordination chemistry, chelators, zirconium, bioconjugation, medical imaging, synucleinopathies

Context: Recent achievements in the synthesis and coordination chemistry of bispidines (bispidine = 3,7-diazabicyclo[3.3.1]nonane) have paved the way to a new field of application of bispidine coordination complexes: medical imaging and more particularly magnetic resonance imaging (MRI)¹ and positron emission tomography (PET).^{2,3} For each of these two techniques, a metal ion (Mn for MRI or a radioisotope for PET) plays the main role of "reporter" and induces the signal. Nevertheless, in both cases, the ions must be chelated and form complexes resistant to *in vivo* dissociation. The particularity of bispidine ligands lies in their rigid and pre-organized skeleton, which allows for fast complexation kinetic in highly dilute conditions suitable for PET and for a strong kinetic inertness of the complexes favourable to their *in vivo* application.

Objective: This project aims at developing bispidine derivatives for the *in vivo* detection of α -synuclein aggregates by PET with ^{89}Zr . Such aggregates are very promising diagnostic and therapeutic targets for synucleinopathies such as Parkinson and Lewy bodies diseases.⁴ We aim at providing a library of new chelates suitable for Zr^{4+} complexation. Their complexation with Zr^{4+} will be characterised by state-of-the art physicochemical techniques (coll. Dr Mourad Elhabiri, CBM/LIMA UMR 7042). The most promising chelates will be conjugated with biomolecules (mAbs, nanobodies) targeted to the synuclein aggregates. Radiolabelling (coll. Dr Frédéric Boisson, IM/UMR7178) and preliminary *in vitro* and *in vivo* imaging will be explored on murine models with diffuse synuclein aggregates and cognitive deficiency (coll. Dr Laurette Boutillier, LNCA).

Expected skills: Multi-stage organic synthesis and standard characterization techniques (NMR, IR, mass spectrometry); Skills and a taste for coordination and physical chemistry; Control of bibliographical research tools; English; Energy, rigour, method and tenacity.

Application/further information: Just contact me at aline.nonat@unistra.fr.




Fig. 1. Bispidine chelators for synuclein imaging: a compromise between fast complexation and strong kinetic inertia.

¹ Ndiaye, D.; Sy, M.; Pallier, A.; Mème, S.; de Silva, I.; Lacerda, S.; Nonat, A. M.; Charbonnière, L. J.; Tóth, E., *Angew. Int. Ed.*, **2020**, *13*, 11958-11963 (cover article).

² Gillet, R.; Roux, A.; Brandel, J.; Huclier-Markai, S.; Camerel, F.; Jeannin, O.; Nonat, A. M.; Charbonnière, L. J., *Inorg. Chem.*, **2017**, *56*, 11738-11752.

³ Price, T. W.; Yap, S.; Gillet, R.; Savoie, H.; Charbonnière, L. J.; Boyle, R.; Nonat, A. M.; Stasiuk, G. J., *Chem. Eur. J.*, **2020**, *18*, 7602-7608 (hot paper).

⁴ M. Colom-Cadena *et al*, *Brain*, **2017**, *140*, 3204–3214.